

# On the Siegel-Weil Theorem for Loop Groups (I)

Howard Garland and Yongchang Zhu\*

## 1 Introduction

The notion of an snt-module and our extension (1.14) of the Siegel-Weil Theorem (also see Theorem 8.1, §8) grew out of our work on the Siegel-Weil Theorem for arithmetic quotients of loop groups, which we prove in Part II of this paper ([3]). In fact (1.14) is a vital step in our proof of the Siegel-Weil theorem in the loop case, and as far as we know, it also seems to be a new result for automorphic forms on certain, finite-dimensional, non-reductive groups. At the same time, the theory of automorphic forms on arithmetic quotients of loop groups, specifically a loop version of Godement's criterion (Theorem 5.3, Part II [3]) is used to prove the convergence theorem (Theorem 3.3) we need for the Eisenstein series  $E_t$  defined in (3.10). At the moment, even though the statement of Theorem 3.3 only involves finite-dimensional groups, the proof using loop groups is the only one we have!

To state our main result of this part, we first recall the Siegel-Weil theorem proved in [7].

Let  $F$  be a number field,  $\mathbf{A}$  be the ring of adeles of  $F$ , and  $M$  be a symplectic space over  $F$  with symplectic pairing  $\langle, \rangle$ . Let  $(V, (, ))$  be a finite dimensional vector space over  $F$  with the non-degenerate, symmetric, bilinear form  $(, )$ . The space  $M \otimes_F V$  has the symplectic form given by

$$\langle u_1 \otimes v_1, u_2 \otimes v_2 \rangle = \langle u_1, u_2 \rangle (v_1, v_2).$$

The group  $Sp_M$  and the orthogonal group  $G$  of  $V$  form a dual pair in the symplectic group  $Sp_{2N}$  of  $M \otimes_F V$  (where  $2N = \dim M \dim V$ ). Let

$$M = M_- \oplus M_+ \tag{1.1}$$

---

\*The second author's research is supported by Hong Kong Research Grant Council earmark grant number HKUST 604507

be a direct sum of Lagrangian subspaces of  $M$ , then

$$M \otimes V = M_- \otimes V \oplus M_+ \otimes V$$

is a direct sum of Lagrangian subspaces of  $M \otimes V$ . The Hilbert space  $L^2((M_- \otimes V)_{\mathbf{A}})$  is an irreducible unitary representation of the metaplectic group  $\widehat{Sp}_{2N}(\mathbf{A})$ , which is called the Weil representation. The dense subspace  $\mathcal{S}((M_- \otimes V)_{\mathbf{A}})$ , formed by the Schwartz-Bruhat functions on  $(M_- \otimes V)_{\mathbf{A}}$ , is invariant under the action of  $\widehat{Sp}_{2N}(\mathbf{A})$ . The theta functional

$$\theta : \mathcal{S}((M_- \otimes V)_{\mathbf{A}}) \rightarrow \mathbb{C}$$

given by

$$\theta(\phi) = \sum_{r \in M_- \otimes V} \phi(r)$$

is  $Sp_{2N}(F)$ -invariant.

The group  $\widehat{Sp}_{2N}(\mathbf{A})$  contains the commuting pair of groups  $\widehat{Sp}_M(\mathbf{A})$  and  $G(\mathbf{A})$ , where  $\widehat{Sp}_M(\mathbf{A})$  is the preimage of  $Sp_M(\mathbf{A})$  in  $\widehat{Sp}_{2N}(\mathbf{A})$ .

For a given  $\phi \in \mathcal{S}((M_- \otimes V)_{\mathbf{A}})$ , there are two simple ways to construct a function on  $Sp_M(F) \backslash \widehat{Sp}_M(\mathbf{A})$ : one is

$$I(\phi, g) \stackrel{\text{def}}{=} \int_{G(F) \backslash G(\mathbf{A})} \theta(\pi(g, h)\phi) dh, \quad g \in \widehat{Sp}_M(\mathbf{A}), \quad (1.2)$$

where  $\pi$  is the Weil representation and  $dh$  is the Haar measure on  $G(\mathbf{A})$  such that the volume of  $G(F) \backslash G(\mathbf{A})$  is 1. For the second way, we first consider the function  $g \mapsto (\pi(g)\phi)(0)$ , which is  $P(\mathbf{A})$ -invariant (where  $P$  is the Siegel parabolic subgroup of  $Sp_M$  that fixes  $M_+$ ). We then form the Eisenstein series

$$E(\phi, g) \stackrel{\text{def}}{=} \sum_{r \in P(F) \backslash Sp_M(F)} (rg \cdot \phi)(0). \quad (1.3)$$

This gives a function on  $Sp_M(F) \backslash \widehat{Sp}_M(\mathbf{A})$ . When  $\dim V > \dim M + 2$ , both (1.2) and (1.3) converge, and the Siegel-Weil formula asserts that the above two constructions are equal, i.e.

$$I(\phi, g) = E(\phi, g). \quad (1.4)$$

Weil [7] proved such an identity in more generality for dual pairs constructed from semisimple algebras with involutions. Generalizations of this formula to non-convergent cases can be found in [5].

Put  $I(\phi) \stackrel{\text{def}}{=} I(\phi, e)$  and  $E(\phi) \stackrel{\text{def}}{=} E(\phi, e)$ , we have

$$I(\phi) = E(\phi) \quad (1.5)$$

Since  $I(\phi, g) = I(g \cdot \phi)$  and  $E(\phi, g) = E(g \cdot \phi)$ , we see that (1.4) and (1.5) are equivalent.

Since the cosets  $P(F) \backslash Sp_M(F)$  are in one-to-one correspondence with the set  $Gr(M)$  of Lagrangian subspaces of  $M$ , via the map  $Pg \mapsto M_+g$  (we assume the symplectic group acts on  $M$  from the right). The Eisenstein series (1.3) can also be written as a summation over  $Gr(M)$  as follows. Let  $\pi_- : M \rightarrow M_-$  denote the projection with respect to the decomposition (1.1). For  $r \in P(F) \backslash Sp_M(F)$ , let  $U = M_+r$ . Then the symplectic pairing  $\pi_-(U) \times M_+ \rightarrow F$  factors through a non-degenerate pairing  $\pi_-(U) \times M_+/(M_+ \cap U) \rightarrow F$ . For each  $v \in \pi_-(U)$ , let  $\tilde{v} \in U$  be a lifting of  $v$ , write  $\tilde{v} = \tilde{v}_- + \tilde{v}_+$  according the decomposition (1.1), then the map

$$\rho : \pi_-(U) \rightarrow M_+/(M_+ \cap U), \quad v \mapsto \tilde{v}_+ + M_+ \cap U \quad (1.6)$$

is well-defined. One can prove that

$$\pi(r)\phi(0) = \int_{(\pi_-(U) \otimes V)_{\mathbf{A}}} \psi\left(\frac{1}{2}\langle x, \rho x \rangle\right) \phi(x) dx \stackrel{\text{def}}{=} E(\phi, U), \quad (1.7)$$

where  $\psi$  is the additive character of  $\mathbf{A}/F$  used in the definition of the Weil representation, and the measure on the right hand side is the Haar measure on  $(\pi_-(U) \otimes V)_{\mathbf{A}}$  normalized by the condition that the covolume of the lattice  $\pi_-(U) \otimes V$  is 1. So the Eisenstein series (1.3) can be written as

$$E(\phi) = \sum_{U \in Gr(M)} E(\phi, U).$$

In our generalization of the Siegel-Weil formula (1.5), we assume the symplectic space  $M$  has an additional structure, which we call an *snt-module*. The groups involved are no longer reductive groups (examples:  $Sp_n(F[t]/(t^k))$  and  $G(F[t]/(t^k))$ ). To state our generalization, we define

**Definition 1.1** *By a symplectic, nilpotent  $t$ -module (=snt-module)  $M$ , we mean an  $F[t]$ -module which is finite-dimensional when considered as a vector space over  $F$ , and which is equipped a symplectic form  $\langle, \rangle$  such that the*

following conditions are satisfied

(i) there exists a positive integer  $N > 0$  such that

$$t^N \cdot \xi = 0, \text{ for all } \xi \in M.$$

(ii). The operator  $t$  is self-dual, i.e.

$$\langle t\xi, \eta \rangle = \langle \xi, t\eta \rangle \quad (1.8)$$

Since  $t^N = 0$  on  $M$ , we may regard  $M$  as an  $F[[t]]$ -module. We give a simple example of an snt-module. Consider  $F[[t]]$ -module

$$H_k = F[[t]]/(t^k) \oplus F[[t]]/(t^k), \quad (1.9)$$

with a symplectic form  $\langle, \rangle$  defined by the conditions

(i). Each of the two summands is isotropic.

(ii).

$$\langle (t^i, 0), (0, t^j) \rangle = \begin{cases} 0, & i + j \neq k - 1 \\ 1, & i + j = k - 1 \end{cases}, \text{ where } i, j = 0, 1, \dots, k - 1.$$

We shall prove later (see Lemma 2.2) that every snt-module is a direct sum of the above examples. For a given snt-module  $M$ ,  $g \in GL(M)$  is called an snt-module automorphism if  $g$  preserves both the  $F[[t]]$ -module structure and the symplectic structure. We denote by

$$Sp(M, t)$$

the group of all snt-automorphisms on  $M$ .

For a given snt-module  $M$  and a space  $V$  with non-degenerate, bilinear, symmetric form  $(, )$ , then the space

$$M \otimes_F V = M \otimes_{F[[t]]} V[[t]] \quad (1.10)$$

has a natural snt-module structure, where  $V[[t]] = V \otimes_F F[[t]]$ . And the symplectic form is defined using the first tensor product:

$$\langle x_1 \otimes v_1, x_2 \otimes v_2 \rangle = \langle x_1, x_2 \rangle (v_1, v_2)$$

and the  $F[[t]]$ -module structure is defined using the second tensor product. Let  $Sp_{2N}$  ( $2N = \dim M \dim V$ ) denote the symplectic group of the symplectic space  $M \otimes_F V$ , the group  $Sp(M, t)$  is a subgroup of  $Sp_{2N}$ . The

orthogonal group  $G(F)$  acts on  $M \otimes_F V$  preserving the snt-module structure. But a larger group  $G(F[[t]])$  acts on (1.10): for  $x \otimes v \in M \otimes_{F[[t]]} V[[t]]$ ,  $g \in G(F[[t]])$ ,

$$(x \otimes v) \cdot g = x \otimes (v \cdot g).$$

It is easy to see that the action preserves both the symplectic structure and the  $F[[t]]$ -module structure, so we also have a group morphism

$$G(F[[t]]) \rightarrow Sp_{2N}.$$

We denote  $G^q(F[[t]])$  the image. The subgroups  $Sp(M, t)$  and  $G^q(F[[t]])$  in  $Sp(M \otimes V, t)$  obviously commute. Our generalization of the Siegel-Weil formula is concerned with the commuting pair  $(Sp(M, t), G^q(F[[t]]))$ , which are not reductive groups in general. Let

$$Gr(M, t)$$

denote the set of all  $F[[t]]$ -stable Lagrangian subspaces of  $M$ , so  $Gr(M, t) \subset Gr(M)$ .

We take a direct sum decomposition  $M = M_- \oplus M_+$  such that  $M_- \in Gr(M)$ ,  $M_+ \in Gr(M, t)$ . As before  $L^2((M_- \otimes V)_{\mathbf{A}})$  is a representation of  $\widehat{Sp}_{2N}(\mathbf{A})$ , with the theta functional

$$\theta : \mathcal{S}((M_- \otimes V)_{\mathbf{A}}) \rightarrow \mathbb{C}, \quad \phi \mapsto \theta(\phi) = \sum_{r \in M_- \otimes V} \phi(r).$$

For a subspace  $U \in Gr(M)$ , let  $E(\phi, U)$  be as in (1.7), and we define

$$\text{Et}(\phi) \stackrel{\text{def}}{=} \sum_{W \in Gr(M, t)} E(\phi, W). \quad (1.11)$$

And we define

$$\text{It}(\phi) \stackrel{\text{def}}{=} \int_{G^q(F[[t]]) \backslash G^q(\mathbf{A}[[t]])} \theta(h \cdot \phi) dh, \quad (1.12)$$

where  $dh$  denotes the Haar measure on  $G^q(\mathbf{A}[[t]])$  such that the volume of  $G^q(F[[t]]) \backslash G^q(\mathbf{A}[[t]])$  is 1.

By Lemma 2.2,  $M$  is isomorphic to a direct sum of  $n$  copies of  $H_k$ 's:

$$M \cong H_{k_1} \oplus \cdots \oplus H_{k_n} \quad (1.13)$$

assume  $\dim V > 6n + 2$ , and the quadratic form  $(\ )$  on  $V$  is  $F$ -anisotropic or  $\dim V - r > \frac{1}{2}\dim M + 1$ , where  $r$  is the dimension of a maximal isotropic

subspace of  $V$ , then we have the following generalization of the Siegel-Weil formula (Theorem 7.3, Theorem 8.1) :

$$\text{Et}(\phi) = \text{It}(\phi). \quad (1.14)$$

The condition  $\dim V > 6n + 2$  is for the convergence of  $\text{Et}(\phi)$ , and the condition that  $(V, (,))$  is  $F$ -anisotropic or  $\dim V - r > \frac{1}{2}\dim M + 1$  is for convergence of  $\text{It}(\phi)$ . This formula reduces to the classical formula (1.5) when  $k_1 = \dots = k_n = 1$ .

In general, the  $Sp(M, t)$ -action on  $Gr(M, t)$  is not transitive, but there are only finitely many orbits, so the sum  $\text{Et}(\phi) = \sum E(U, \phi)$  is a sum of Eisenstein series induced from several "parabolic" subgroups (rather than only one), and each corresponds to a  $Sp(M, t)$ -orbit in  $Gr(M, t)$ .

In the case that  $M$  is a direct sum of  $n$ -copies of the snt-module  $H_k$ , then  $Sp(M, t) = Sp_{2n}(F[t]/(t^k))$  and  $G^q(F[[t]]) = G(F[t]/(t^k))$ . The formula (1.14) means that the Siegel-Weil formula holds for symplectic and orthogonal groups over  $F[t]/(t^k)$ . It implies that the Siegel-Weil formula holds for symplectic and orthogonal groups over  $F[t]/(p(t))$  for arbitrary polynomial  $p(t)$ .

We now give an explicit example of our formula. Let  $F = \mathbb{Q}$ , and the snt - module  $M$  be

$$M = \mathbb{Q}[t]/(t^2) \oplus \mathbb{Q}/(t^2) \quad (1.15)$$

with the snt-module structure given as in (1.9). And we take a positive definite even unimodular lattice  $L$  of rank  $N$  with the bilinear form given by  $(,)$ , and let  $V = \mathbb{Q}L$ . It is well-known that  $N$  is divisible by 8. Let  $L_1, \dots, L_g$  be the list of the positive definite even unimodular lattices of rank  $N$  (up to isomorphism), let  $(,)_j$  denote the pairing of  $L_j$ , and  $|\text{Aut}_j|$  be the order of automorphism group of  $L_j$ . We denote  $1 \oplus 0$  and  $0 \oplus 1$  in (1.15) by  $e_1$  and  $e_2$  respectively, then  $e_1, te_1, e_2, te_2$  is a  $\mathbb{Q}$ -basis of  $M$ . Let  $M_- = \mathbb{Q}e_1 + \mathbb{Q}te_1$  and  $M_+ = \mathbb{Q}te_2 + \mathbb{Q}e_2$ . It is clear that  $M_-$  and  $M_+$  are Lagrangian subspaces of  $M$  and

$$M = M_- \oplus M_+.$$

We have the Weil representation of  $Sp_{4N}(\mathbf{A})$  on

$$\mathcal{S}((M_- \otimes V)_{\mathbf{A}}).$$

Take  $\phi = \Pi_v \phi_v \in \mathcal{S}((M_- \otimes V)_{\mathbf{A}})$  as follows: for a finite place  $p$  of  $\mathbb{Q}$ ,  $\phi_p$  is the characteristic function of  $e_1 \otimes L_{\mathbf{Z}_p} + te_1 \otimes L_{\mathbf{Z}_p}$  (where  $\mathbf{Z}_p$  denotes the

ring of  $p$ -adic integers and  $L_{\mathbf{Z}_p} = L \otimes \mathbf{Z}_p$ ); for the real place  $\infty$  of  $\mathbb{Q}$ ,

$$\phi_\infty(e_1 \otimes v_1 + te_1 \otimes v_2) = e^{\pi i \tau_1(v_1, v_1) + \pi i \frac{-1}{\tau_2}(v_2, v_2)},$$

where  $\tau_1, \tau_2$  are complex numbers in the upper half plane. With the above choice of  $\phi$ , our new Siegel-Weil formula (1.14) becomes

$$\begin{aligned} & 1 + \frac{1}{2} \sum_{a \geq 1, b \in \mathbf{Z}: (a, b) = 1} \sum_{m, n \in \mathbf{Z}: (m, n) = 1} (am^2\tau_1 + an^2\tau_2 + b)^{-\frac{N}{2}} \\ &= C \sum_{j=1}^g \frac{1}{|\text{Aut}_j|} \sum_{u, v \in L_j, u, v \text{ colinear}} e^{\pi i \tau_1(u, u) + \pi i \tau_2(v, v)} \end{aligned} \quad (1.16)$$

where  $(m, n) = 1$  ( $(a, b) = 1$ ) means  $m, n$  (resp.  $a, b$ ) are relatively prime, and  $u, v$  colinear means the  $\mathbb{Q}$ -span of  $u, v$  is at most 1-dimensional. And the constant  $C$  is given by (1.18) below.

We compare (1.16) with the classical Siegel-Weil formula

$$\frac{1}{2} \sum_{m, n \in \mathbf{Z}, (m, n) = 1} (m\tau + n)^{-\frac{N}{2}} = C \sum_{j=1}^g \frac{1}{|\text{Aut}_j|} \sum_{u \in L_j} e^{\pi i \tau(u, u)} \quad (1.17)$$

This formula expresses the  $\frac{N}{2}$ -th Eisenstein series of  $SL(2, \mathbb{Z})$  as a sum of Theta series. The constant term in  $q$ -expansion of both sides gives the density formula

$$1 = C \sum_{j=1}^g \frac{1}{|\text{Aut}_j|}, \quad (1.18)$$

which determines  $C$ .

If we take more general test function  $\phi$ , our Siegel-Weil formula (1.14) is

$$\begin{aligned} & 1 + \frac{1}{2} \sum_{a \geq 1, b \in \mathbf{Z}: (a, b) = 1} \sum_{m, n \in \mathbf{Z}: (m, n) = 1} (am^2\tau_{11} + 2amn\tau_{12} + an^2\tau_{22} + b)^{-\frac{N}{2}} \\ &= C \sum_{j=1}^g \frac{1}{|\text{Aut}_j|} \sum_{u, v \in L_j, u, v \text{ colinear}} e^{\pi i (\tau_{11}(u, u) + 2\tau_{12}(u, v) + \tau_{22}(v, v))}, \end{aligned} \quad (1.19)$$

where  $C$  is determined by (1.18).

The paper is organized as follows: In §2 we study the structure of an snt-module  $M$  (Lemma 2.2) and the structure of the corresponding group  $Sp(M, t)$  of snt-automorphisms of  $M$  (Cor. 2.6). In §3 we recall basic facts about the Weil representation and study the Eisenstein series  $Et(\phi)$  for the Weil representation associated to an snt-module. In particular, we state the convergence theorem (Theorem 3.3) for such Eisenstein series. As we already stated, the proof depends on the convergence theorem for Eisenstein series on loop groups, and will be given in Part II of this paper ([3]). We also state a consequence, Theorem 3.4, of Theorem 3.3 together with Prop. 1 and Prop. 2 of [7].

In §4 we extend a result in Weil [7] and obtain Theorem 4.7, which identifies certain abstract measures associated with the map  $T_W$  (see (3.13) for the definition of  $T_W$ ) with certain gauge measures in the sense of [7]. In §5 we obtain Theorems 5.4 and 5.8. In particular, we identify the space of  $G(F[[t]])$  - orbits in  $M_- \otimes V$  with the set of pairs  $(W, i)$  with  $W \in Gr(M_-, t)$  (the  $t$  - Grassmannian),  $i \in S_t^2(W)$  such that  $U(i)_F$  is non-empty (  $U(i)_F$  is defined in §5, just before the statement of Theorem 5.8).

In §6 we discuss  $\theta$  - series and finally, we prove Theorems 7.3 and 8.1, our versions of the Siegel-Weil theorem for snt-modules.

## 2 The structure of symplectic, nilpotent $t$ -modules.

In this section, we prove some results about the structure of symplectic, nilpotent  $t$ -modules (snt-modules) defined in Section 1 (Definition 1.1).

**Lemma 2.1** *Let  $M$  be an snt-module. For all  $\xi \in M$ ,  $k \in \mathbb{Z}_{>0}$ , we have*

$$\langle \xi, t^k \xi \rangle = 0.$$

*Proof.* We have for all  $\xi \in M$ ,  $k \in \mathbb{Z}_{\geq 0}$ ,

$$\langle \xi, t^k \cdot \xi \rangle = -\langle t^k \cdot \xi, \xi \rangle,$$

since  $\langle \cdot, \cdot \rangle$  is skew symmetric. On the other hand

$$\langle \xi, t^k \cdot \xi \rangle = \langle t^k \cdot \xi, \xi \rangle$$

by (1.8). Hence

$$\langle \xi, t^k \cdot \xi \rangle = 0,$$

□



**Lemma 2.2** *Every snt-module  $M$  is isomorphic to a direct sum*

$$M \cong H_{k_1} \oplus \dots \oplus H_{k_n}$$

$k_1 \geq k_2 \geq \dots \geq k_n$ . Where  $H_k$  is given as (1.9). Moreover,  $n$  and the  $k_i$  are uniquely determined by  $M$ .

*Proof.* The uniqueness is clear from the theory of elementary divisors. For  $\xi \in M$ , we define the order of  $\xi$  to be the smallest positive integer  $k$ , such that  $t^k \cdot \xi = 0$ . Pick  $\xi \in M$  of maximal order ( $N$  say). Then the vectors

$$\xi, t \cdot \xi, \dots, t^{N-1} \cdot \xi$$

are linearly independent over  $F$ . To see this, we consider  $F[[t]]$ -submodule  $F[[t]]\xi$ . Since  $F[[t]]$  is a PID,  $F[[t]]\xi$  is isomorphic to  $F[[t]]/(t^k)$ . It is clear that  $k = N$ .

Since  $\langle, \rangle$  is non-degenerate, we can find  $\eta \in M$ , so that

$$\langle t^{N-1} \cdot \xi, \eta \rangle = 1, \quad \langle t^j \cdot \xi, \eta \rangle = 0, \quad j = 0, 1, \dots, N-2$$

But then,

$$\langle \xi, t^{N-1} \cdot \eta \rangle = \langle t^{N-1} \cdot \xi, \eta \rangle = 1$$

Hence

$$t^{N-1} \cdot \eta \neq 0,$$

and so

$$\eta, t \cdot \eta, \dots, t^{N-1} \cdot \eta$$

are linearly independent (by the above argument) and

$$t^N \cdot \eta = 0$$

(since  $N$  was assumed the maximal order of any element of  $M$ ). But then if

$$C_1 = F\text{-span of } \xi, t \cdot \xi, \dots, t^{N-1} \cdot \xi,$$

$$C_2 = F\text{-span of } \eta, t \cdot \eta, \dots, t^{N-1} \cdot \eta,$$

we see that

$$H = C_1 \oplus C_2$$

with symplectic structure given by the restriction of  $\langle, \rangle$  on  $M$  is isomorphic to the snt-module  $H_N$ , with  $C_1, C_2$  corresponding to the two direct

summands  $F[[t]]/(t^N)$ . Since  $\langle, \rangle$  restricted to  $H$  is non-degenerate,  $M$  decomposes as a direct sum of *snt*-modules

$$M \cong H \oplus H^\perp,$$

and applying the induction hypothesis to  $H^\perp$ , we obtain the lemma.  $\square$

For an *snt*-module  $M$ , so  $M$  is in particular a symplectic space. As in Section 1, we let  $Gr(M)$  denote the set of all Lagrangian subspaces, and  $Gr(M, t)$  denote the set of Lagrangian subspaces which are stable under the action of  $F[[t]]$ , so  $Gr(M, t) \subset Gr(M)$ . We call elements in  $Gr(M, t)$  *t*-Lagrangian subspaces. We have:

**Lemma 2.3** *If  $U \subset M$  is a  $F[[t]]$ -stable, isotropic and it is not properly contained in any other  $F[[t]]$ -stable, isotropic subspaces, then  $U$  is *t*-Lagrangian, i.e.  $U \in Gr(M, t)$ .*

*Proof.* Assume  $U$  is not maximal isotropic. Then there exists  $v \in M$ ,  $v \notin U$ , such that

$$\langle v, u \rangle = 0, \quad \text{all } u \in U.$$

But then for all  $u \in U$ ,  $i \in \mathbb{Z}_{\geq 0}$

$$\langle t^i \cdot v, u \rangle = \langle v, t^i \cdot u \rangle = 0,$$

since  $t^i \cdot u \in U$ .

Hence, the space  $Span_{F[[t]]}\{v, U\}$  is  $F[[t]]$ -stable, isotropic and contains  $M$  properly. This contradicts the maximality of  $M$ .  $\square$

As in Section 1, we let  $Sp(M, t)$  be the subgroup of all  $\sigma \in Sp(M)$  such that  $\sigma$  is an  $F[[t]]$ -module automorphism. For a finite dimensional  $F[[t]]$ -module  $U$ , a subset of non-zero elements  $e_1, \dots, e_n$  is called a *quasi-basis* of  $U$  if every element in  $U$  can be written as an  $F[[t]]$ -linear combination of  $e_i$ 's and  $a_1 e_1 + \dots + a_n e_n = 0$  implies that all  $a_i e_i = 0$ . For example, for

$$U = F[[t]]/(t^{k_1}) \oplus \dots \oplus F[[t]]/(t^{k_n}),$$

the set (n elements)  $(1, 0, \dots, 0), (0, 1, \dots, 0), \dots, (0, \dots, 1)$  is a quasi-basis. If  $U$  has a quasi-basis consisting of  $n$ -elements, then the  $F$ -vector space  $\bar{U} \stackrel{\text{def}}{=} U/tU$  is  $n$ -dimensional. If we have two such  $F[[t]]$ -modules  $U_1$  and  $U_2$ , and  $T : U_1 \rightarrow U_2$  is a morphism, then  $T$  induces a  $F$ -linear map  $\bar{T} : \bar{U}_1 \rightarrow \bar{U}_2$ .

An *snt*-module  $M$  with decomposition as Lemma 2.2 is called *homogeneous* if  $k_1 = k_2 = \dots = k_n = k$ . In this case the group  $Sp(M, t)$  is determined by

**Lemma 2.4** *Let  $M$  be a homogeneous snt-module as in (1.13) with  $k_1 = \dots = k_n = k$ . Then  $Sp(M, t)$  is isomorphic to  $Sp_{2n}(F[[t]]/(t^k))$ . In particular, its reduction mod  $t$  defines a surjective group homomorphism*

$$\pi_0 : Sp(M, t) \rightarrow Sp_{2n}(F),$$

*Proof.* We first consider the free  $F[[t]]/(t^k)$ -module  $(F[[t]]/(t^k))^{2n}$ . It has as standard  $F[[t]]/(t^k)$ -valued symplectic form  $\langle, \rangle^\wedge$ . We define an  $F$ -valued symplectic form  $\langle, \rangle$  on  $(F[[t]]/(t^k))^{2n}$  by

$$\langle a, b \rangle = \text{coefficient of } t^{k-1} \text{ in } \langle a, b \rangle^\wedge.$$

With this  $\langle, \rangle$ ,  $(F[[t]]/(t^k))^{2n}$  is an snt-module, which is clearly isomorphic to  $M$  in the lemma. It follows from the construction that  $Sp_{2n}(F[[t]]/(t^k))$  preserves the snt-module structure, so we have

$$Sp_{2n}(F[[t]]/(t^k)) \subset Sp(M, t).$$

For the converse inclusion is also clear. □

For a homogeneous snt-module  $M$  as in Lemma 2.4,  $\bar{M} = M/tM$  has a symplectic structure defined as follows, if  $\bar{a}, \bar{b} \in \bar{M}$ , let  $a, b \in M$  be their liftings, then

$$\langle \bar{a}, \bar{b} \rangle \stackrel{\text{def}}{=} \langle a, t^{k-1}b \rangle. \quad (2.1)$$

Now we turn to the case of general (possibly non-homogeneous) snt-modules  $M$ , with direct sum decomposition as in Lemma 2.2. For fixed  $k$ , we let

$$M(k) = \oplus_{k_i=k} H_{k_i},$$

so

$$M = M(l_1) \oplus \dots \oplus M(l_s), \quad l_1 > l_2 > \dots > l_s, \quad (2.2)$$

with each  $M(l_i)$  a homogeneous snt-submodule. The  $M(l_i)$ 's are mutually orthogonal with respect to  $\langle, \rangle$ .

Now if  $\sigma \in Sp(M, t)$ , then  $\sigma$  (acting on the right, recall) has a block decomposition with respect to (2.2),

$$\sigma = \begin{bmatrix} \sigma_1^1 \dots \sigma_s^1 \\ \dots \\ \sigma_1^s \dots \sigma_s^s \end{bmatrix} \quad (2.3)$$

where  $\sigma_j^i : M(l_i) \rightarrow M(l_j)$ , and for  $\xi = (\xi_1, \dots, \xi_s) \in M$  ( $\xi_i \in M(l_i)$ )

$$\xi\sigma = (\xi_1, \dots, \xi_s) \begin{bmatrix} \sigma_1^1 \dots \sigma_s^1 \\ \dots \\ \sigma_1^s \dots \sigma_s^s \end{bmatrix}.$$

We have also the decomposition  $\bar{M} = M/tM$  induced from the decomposition (2.2),

$$\bar{M} = \bar{M}(l_1) \oplus \dots \oplus \bar{M}(l_s), \quad l_1 > l_2 > \dots > l_s, \quad (2.4)$$

So  $\bar{\sigma} : \bar{X} \rightarrow \bar{X}$  has a block decomposition:

$$\bar{\sigma} = \begin{bmatrix} \bar{\sigma}_1^1 \dots \bar{\sigma}_s^1 \\ \dots \\ \bar{\sigma}_1^s \dots \bar{\sigma}_s^s \end{bmatrix} \quad (2.5)$$

**Lemma 2.5** *The matrix  $\bar{\sigma}$  is block-upper triangular, i.e.*

$$\bar{\sigma}_i^j = 0 \text{ for } i < j \quad (2.6)$$

and the diagonal block  $\bar{\sigma}_i^i$  is in  $Sp(\bar{X}(l_i))$  (recall  $\bar{X}(l_i)$  has the symplectic structure defined by (2.1)).

*Proof.* For  $i < j$ ,  $v_j \in M(l_j)$ , we have  $t^{l_j}v_j = 0$  and  $\sigma_i^j$  is  $t$ -linear, this implies that  $t^i(v_j\sigma_i^j) = 0$ , then

$$v_j\sigma_i^j \in t^{l_i-l_j}M(l_i) \quad (2.7)$$

This implies that  $\bar{\sigma}_i^j = 0$ . For  $a, b \in M(l_i)$ , we have

$$\langle t^{l_i-1}a, b \rangle = \langle t^{l_i-1}a\sigma, b\sigma \rangle = \sum_{j=1}^s \langle t^{l_i-1}a\sigma_j^i, b\sigma_j^i \rangle, \quad (2.8)$$

if  $j < i$ , by (2.7) we have

$$t^{l_i-1}a\sigma_j^i \in t^{l_i-1}t^{l_j-l_i}M(l_j) = t^{l_j-1}M(l_j)$$

and

$$b\sigma_j^i \in t^{l_j-l_i}M(l_j)$$

it implies that

$$\langle t^{l_i-1}a\sigma_j^i, b\sigma_j^i \rangle = 0.$$

For  $j > i$ , then  $l_i - 1 \geq l_j$ , we have

$$t^{l_i-1}a\sigma_j^i \in t^{l_i-1}M(l_j) = 0.$$

So (2.8) gives that

$$\langle t^{l_i-1}a, b \rangle = \langle t^{l_i-1}a\sigma_i^i, b\sigma_i^i \rangle,$$

by the definition (2.1) of the symplectic form on  $\bar{M}(l_i)$ , we prove that  $\bar{\sigma}_i^i$  is a symplectic isomorphism of  $\bar{M}(l_i)$ .  $\square$

**Corollary 2.6** *Let  $M$  be an snt-module as in (2.2), then  $Sp(M, t)$  is the semi-direct product*

$$Sp(M, t) = N \ltimes H,$$

where  $N$  is the unipotent radical of  $Sp(M, t)$  and

$$H \cong \prod_{i=1}^s Sp_{2r_i}(F),$$

where  $r_i$  is the number  $H_{l_i}$ 's in the decomposition of  $M(l_i)$ .

Next we discuss the classification of  $t$ -Lagrangian subspaces. First for the snt-module (1.9), let  $e_1, e_2$  denote  $(1, 0), (0, 1)$  respectively. For each  $0 \leq i \leq k-1$ , let  $L_i$  be the  $F$ -subspace with basis

$$t^i e_1, t^{i+1} e_1, \dots, t^{k-1} e_1, t^{k-i} e_2, t^{k-i+1} e_2, \dots, t^{k-1} e_2.$$

It is clear that  $L_i$  is a  $t$ -Lagrangian subspace. For an snt-module  $M$  with decomposition as in Lemma 2.2, let  $L_{i_j} \subset H_{k_j}$  be the subspace described as above, it is clear that

$$L_{i_1} \oplus \dots \oplus L_{i_n} \tag{2.9}$$

is an  $t$ -Lagrangian subspace of  $M$ . We have

**Proposition 2.7** *Let  $M$  be a snt-module with decomposition as in Lemma 2.2, then every  $t$ -Lagrangian subspace can be transformed by some  $g \in Sp(M, t)$  to an  $t$ -Lagrangian subspace as in (2.9).*

This proposition will not be used later, we skip its proof.

### 3 The Weil representation and $t$ -Eisenstein series.

In this section, we first recall some basic facts about the Weil representation associated to a symplectic space over  $F$ , then we study the Eisenstein series  $\text{Et}(\phi)$  (1.11) for the Weil representations associated to an snt-module.

We shall fix a non-trivial additive character  $\psi : \mathbf{A} \rightarrow S^1$  that is trivial on  $F$ . Let  $F^{2N}$  be the standard symplectic space over  $F$ , and  $C = C_- \oplus C_+$  be a direct sum into Lagrangian subspaces; then a two-fold cover, denoted by  $\widehat{Sp}_{2N}(\mathbf{A})$ , of the adelic group  $Sp_{2N}(\mathbf{A})$  acts on  $L^2(C_{-,\mathbf{A}})$ . The subspace  $\mathcal{S}(C_{-,\mathbf{A}})$ , formed by the Schwartz-Bruhat functions is invariant under this action. We recall now the action formula. For  $g \in Sp_{2N}(\mathbf{A})$ , let

$$\begin{bmatrix} \alpha_g & \beta_g \\ \gamma_g & \delta_g \end{bmatrix} \quad (3.1)$$

be the block decomposition of  $g$  with respect to the decomposition

$$\mathbf{A}^{2N} = C_{-,\mathbf{A}} \oplus C_{+,\mathbf{A}}.$$

In this paper, we always assume the action of  $Sp_{2N}$  on  $F^{2N}$  (as well as other symplectic group actions on symplectic spaces) is from the right. So  $\gamma_g$  in (3.1) is a map from  $C_+ \rightarrow C_-$ . Let  $\tilde{g} \in \widehat{Sp}_{2N}(\mathbf{A})$  be a lifting of  $g$ . For  $\phi \in \mathcal{S}(C_{-,\mathbf{A}})$ ,  $(\tilde{g} \cdot \phi)(x)$  equals to

$$\lambda \int_{\text{Im } \gamma_g} S_g(x + x^*) \phi(x\alpha_g + x^*\gamma_g) d(x^*\gamma_g), \quad (3.2)$$

where  $\lambda \in \mathbb{C}^*$  is a certain scalar depending only on  $\tilde{g}$ ,  $d(x^*\gamma_g)$  is a Haar measure on  $\text{Im } \gamma_g$  and

$$S_g(x + x^*) = \psi \left( \frac{1}{2} \langle x\alpha_g, x\beta_g \rangle + \frac{1}{2} \langle x^*\gamma_g, x^*\delta_g \rangle + \langle x^*\gamma_g, x\beta_g \rangle \right);$$

it is easy to see that  $\langle x^*\gamma_g, x^*\delta_g \rangle$  depends only on  $x^*\gamma_g$  (not on the choice of  $x^*$ ), therefore  $S_g(x + x^*)$  is a function of  $x$  and  $x^*\gamma_g$ . Since  $\tilde{g}$  is unitary,  $\lambda$  can be determined up to a factor in  $S^1$ .

Let  $P$  be the subgroup of  $Sp_{2N}$  that consists of elements that maps  $C_+$  to itself. An element  $g$  is in  $P(\mathbf{A})$  iff  $\gamma_g = 0$ . Then there is a lifting  $P(\mathbf{A}) \subset \widehat{Sp}_{2N}(\mathbf{A})$  so that for  $g \in P(\mathbf{A})$  and  $\phi \in \mathcal{S}(C_{-,\mathbf{A}})$ ,

$$(g \cdot \phi)(x) = |\det(\alpha_g)|_{\mathbf{A}}^{\frac{1}{2}} \psi\left(\frac{1}{2} \langle x\alpha_g, x\beta_g \rangle\right) \phi(x\alpha_g), \quad (3.3)$$

where the factor  $|det(\alpha_g)|_{\mathbf{A}}^{\frac{1}{2}}$  guarantees the unitarity of the operator  $g$ . There is also a lifting

$$Sp_{2N}(F) \subset \widehat{Sp}_{2N}(\mathbf{A})$$

such that theta functional

$$\theta : \mathcal{S}(C_{-, \mathbf{A}}) \rightarrow \mathbb{C}$$

given by

$$\theta(\phi) = \sum_{r \in C_-} \phi(r)$$

is invariant under  $Sp_{2N}(F)$ . The action of  $Sp_{2N}(F)$  is given by (3.2) with  $\lambda = 1$  and the Haar measure is given by the condition that the covolume of  $(\text{Im } \gamma_g)(F)$  is 1.

For a given snt-module  $M$  with

$$M = H_{k_1} \oplus \cdots \oplus H_{k_n} \quad (3.4)$$

with  $H_{k_i}$  is as in (1.9), and a space  $V$  with non-degenerate, bilinear, symmetric form  $(, )$ . Let  $G$  denote the orthogonal group of  $V$ . Recall that

$$M \otimes_F V = M \otimes_{F[[t]]} V[[t]] \quad (3.5)$$

has a natural snt-module structure (Section 1). Let  $Sp_{2N}$  (where  $2N = \dim M \dim V$ ) denote the symplectic group of the symplectic space  $M \otimes_F V$ . The group  $Sp(M, t)$  is a subgroup of  $Sp_{2N}$ . The group  $G(F[[t]])$  acts on  $M \otimes_{F[[t]]} V[[t]]$  in the second factor, so we have a group morphism

$$G(F[[t]]) \rightarrow Sp(M \otimes V, t).$$

Suppose  $l = \max(k_1, \dots, k_n)$ ; then the image  $G^q(F[[t]])$  of the above homomorphism is isomorphic to  $G(F[[t]]/(t^l))$ . We have a commuting pair  $(Sp(M, t), G(F[[t]]/(t^l)))$  in  $Sp(M \otimes_F V, t)$ .

Suppose we have a direct sum decomposition

$$M = M_- \oplus M_+ \quad (3.6)$$

such that  $M_+, M_- \in Gr(M, t)$ . We put

$$X \stackrel{\text{def}}{=} M_- \otimes V.$$

The space  $L^2(X_{\mathbf{A}})$  is a representation of metaplectic group  $\widehat{Sp}_{2N}(\mathbf{A})$ , and we have the theta functional

$$\theta : \mathcal{S}(X_{\mathbf{A}}) \rightarrow \mathbb{C}, \quad \phi \mapsto \theta(\phi) = \sum_{r \in X} \phi(r). \quad (3.7)$$

Recall the Eisenstein series (1.3), (1.7) for  $\phi \in \mathcal{S}(X_{\mathbf{A}})$  is given by

$$E(\phi) = \sum_{U \in Gr(M)} E(\phi, U) = \sum_{U \in Gr(M)} \int_{(\pi_-(U) \otimes V)_{\mathbf{A}}} \psi\left(\frac{1}{2}\langle x, \rho x \rangle\right) \phi(x) dx.$$

Let  $\pi_- : M \rightarrow M_-$  be the projection map with respect to (3.6). It gives a map

$$Gr(M) \rightarrow Gr(M_-) : \quad U \mapsto \pi_-(U). \quad (3.8)$$

We wish to describe the inverse image of a given  $W \in Gr(M_-)$ . Let

$$W^\perp = \{x \in M_+ \mid \langle W, x \rangle = 0\}.$$

If  $U \in Gr(M)$  satisfying  $\pi_-(U) = W$ , then it is easy to see that

$$W^\perp = M_+ \cap U.$$

The symplectic pairing  $W \times M_+ \rightarrow F$  factors through a non-degenerate pairing

$$\langle \rangle : W \times M_+/W^\perp \rightarrow F.$$

We may identify  $W^*$  with  $M_+/W^\perp$  using this pairing. Recall we have the map

$$\rho_U : W \rightarrow M_+/W^\perp$$

as defined in (1.6). It is easy to prove that  $\rho_U$  is self-dual. We have

**Lemma 3.1** *For a given  $W \in Gr(M_-)$ , the map*

$$U \mapsto \rho_U$$

*is a bijection from the set of  $U \in Gr(M)$  such that  $\pi_-(U) = W$  to the set self-dual linear maps from  $W$  to  $W^* = M_+/W^\perp$*

*Proof.* It is clear that the map  $U \mapsto \rho_U$  is one-to-one. If  $\rho : W \rightarrow W^* = M_+/W^\perp$  is self-dual, then

$$U \stackrel{\text{def}}{=} \{w + \rho(w) + W^\perp \mid w \in W\} \quad (3.9)$$



is a Lagrangian subspace of  $M$ , and  $\rho_U = \rho$ . So the map in the lemma is also onto.  $\square$

Recall the  $t$ -Eisenstein series defined in (1.11) is a sub-series of  $E(\phi)$  given by

$$\text{Et}(\phi) = \sum_{U \in \text{Gr}(X, t)} E(\phi, U). \quad (3.10)$$

Since  $M_-$  and  $M_+$  are  $F[[t]]$ -submodules of  $M$ , the projection map

$$\pi_- : M \rightarrow M_-$$

is an  $F[[t]]$ -module homomorphism. For each  $U \in \text{Gr}(M, t)$ ,  $\pi_-(U)$  is an  $F[[t]]$ -submodule of  $M_-$ . Denote  $\text{Gr}(M_-, t)$  the set of  $F[[t]]$ -submodules of  $M_-$ , so we have map

$$P : \text{Gr}(M, t) \rightarrow \text{Gr}(M_-, t) : U \mapsto \pi_-(U). \quad (3.11)$$

For a  $W \in \text{Gr}(M_-, t)$ , we wish to describe the inverse image

$$P_W \stackrel{\text{def}}{=} P^{-1}(W).$$

For any  $U \in P_W$ , i.e.  $U \in \text{Gr}(M, t)$  and  $\pi_-(U) = W$ , we have the linear map

$$\rho_U : W \rightarrow M_+/W^\perp,$$

as defined in (1.6). As before  $\rho_U$  is self-dual. We now prove  $\rho_U$  is  $F[t]$ -linear. If  $w \in W$ , since  $\pi(U) = W$ , there is  $w' \in W_+$  such that  $w + w' \in U$ . By our definition of  $\rho_U$ ,  $\rho_U(w) = w' \bmod(W^\perp)$ . Since  $U$  is  $t$ -Lagrangian,  $tw + tw' \in U$ , this implies  $\rho_U(tw) = tw' \bmod(W^\perp)$ .

**Lemma 3.2** *For each  $U \in P_W$ ,*

$$\rho_U : W \rightarrow M_+/W^\perp = M_+/(M_+ \cap U)$$

*is  $F[[t]]$ -linear and self-dual. Conversely for each  $\rho : W \rightarrow M_+/W^\perp$  that is  $F[[t]]$ -linear and self-dual, there is a unique  $U \in P_W$  such that  $\rho_U = \rho$ . Therefore  $U \mapsto \rho_U$  is a bijection from  $P_W$  to  $F_W$ , the space of all  $\rho : W \rightarrow M_+/W^\perp$  that is self-dual and  $F[[t]]$ -linear.*

This lemma is an  $t$ -analog of Lemma 3.1. Suppose  $\rho$  is  $F[[t]]$ -linear and self-dual, then  $U$  given as (3.9) is the unique  $t$ -Lagrangian subspace such that  $\rho_U = \rho$ .

We set

$$\text{Et}_W(\phi) = \sum_{U \in P_W} E(\phi, U),$$

By Lemma 3.2, we have

$$\text{Et}_W(\phi) = \sum_{\rho \in F_W} \int_{(W \otimes V)_{\mathbf{A}}} \phi(x) \psi\left(\frac{1}{2} \langle x, \rho(x) \rangle\right) dx, \quad (3.12)$$

where  $dx$  denotes the Haar measure on  $(W \otimes V)_{\mathbf{A}}$  such that the covolume of  $W \otimes V$  is 1. We have

$$\text{Et}(\phi) = \sum_{W \in \text{Gr}(M_-, t)} \text{Et}_W(\phi).$$

For an  $F[[t]]$ -submodule  $W \subset M_-$ , we let

$$S_t^2(W) \subset W \otimes_{F[[t]]} W$$

denote the  $F[[t]]$ -submodule of symmetric tensors. We define

$$T_W : W \otimes_F V = W \otimes_{F[[t]]} V[[t]] \rightarrow S_t^2(W)$$

by

$$T_W : \sum_{i=1}^s w_i \otimes v_i \mapsto \sum_{i,j=1}^s (v_i, v_j) w_i \otimes w_j, \quad (3.13)$$

where  $w_i \in W$ ,  $v_i \in V$ ,  $i = 1, \dots, s$ , and where  $w_i \otimes w_j$  denotes the tensor product of  $w_i, w_j$  in  $S_t^2(W)$ . Of course  $T_W$  can be extended to an adelic map

$$T_W : (W \otimes_F V)_{\mathbf{A}} \rightarrow S_t^2(W)_{\mathbf{A}},$$

and for  $r \in S_t^2(W)_{\mathbf{A}}$ , we set

$$\mathcal{U}_r = \{x \in (W \otimes_F V)_{\mathbf{A}} \mid T_W(x) = r\}.$$

We now consider  $\text{Et}_W(\phi)$  as defined in (3.12). In that expression we consider  $\rho \in F_W$ . We have a pairing

$$S_t^2(W) \times F_W \rightarrow F, \quad \left(\sum w_i \otimes u_i, \rho\right) = \sum \langle w_i, \rho(u_i) \rangle, \quad (3.14)$$

which is clearly non-degenerate. And we have

$$\langle w, \rho(w) \rangle = (T_W(w), \rho), \quad w \in (W \otimes V)_{\mathbf{A}}.$$

We may then rewrite the right hand side of (3.12) as

$$\sum_{\rho \in F_W} \int_{(W \otimes V)_{\mathbf{A}}} \phi(x) \psi\left(\frac{1}{2}(T_W(x), \rho)\right) dx. \quad (3.15)$$

The following result is a corollary of convergence of Eisenstein series on loop groups, it will be proved in part II.

**Theorem 3.3** *Let  $M$  be an  $snt$ -module as in (3.4), where  $H_k$  is as in (1.9). If  $\dim V > 6n+2$ , then the series (3.10) converges absolutely and the convergence is uniform for  $\phi$  varying over a compact subsets in  $\mathcal{S}(X_{\mathbf{A}})$ . It follows that the series (3.12) = (3.15) converges absolutely and the convergence is uniform for  $\phi$  varying over a compact subset of  $\mathcal{S}((W \otimes V)_{\mathbf{A}})$ .*

We can apply Proposition 1 and Proposition 2 of [7]. Using Weil's notation as in [7]:

$$X = (W \otimes V)_{\mathbf{A}}, \quad G = S_t^2(W)_{\mathbf{A}}, \quad \Gamma = S_t^2(W), \quad f = T_W. \quad (3.16)$$

and  $G^* = (F_W)_{\mathbf{A}}$ ,  $\Gamma_* = F_W$ , where we regard  $G^* = (F_W)_{\mathbf{A}}$  as Pontryagin dual of  $G$  by the pairing

$$S_t^2(W)_{\mathbf{A}} \times (F_W)_{\mathbf{A}} \rightarrow S^1, \quad \{a, \rho\} = \psi\left(\frac{1}{2}(a, \rho)\right).$$

We have

$$F_{\phi}^*(g^*) = \int_{(W \otimes V)_{\mathbf{A}}} \phi(x) \psi\left(\frac{1}{2}\{T_W(x), g^*\}\right) dx = \int_X \phi(x) \{f(x), g^*\} dx.$$

Theorem 3.3 implies that the condition of Proposition 2 in [7] is satisfied; that is

$$\sum_{r^* \in \Gamma_*} |F_{\Phi}^*(g^* + \gamma^*)|$$

converges and the convergence is uniform as  $(\phi, g^*)$  varies over a compact subset of  $\mathcal{S}(X) \times G^*$ . By using of Proposition 1 and Proposition 2 [7], we obtain

**Theorem 3.4** Suppose  $\dim V > 6n + 2$ . To every  $r \in S_t^2(W)_{\mathbf{A}}$ , there corresponds a unique positive measure  $\mu_r$  on  $(W \otimes V)_{\mathbf{A}}$  whose support is contained in  $\mathcal{U}_r$ , so that for every function  $\phi$  on  $(W \otimes V)_{\mathbf{A}}$  which is continuous with compact support, the function  $F_\phi(r) = \int \phi d\mu_r$  is continuous and satisfies

$$\int F_\phi(r) dr = \int \phi(x) dx$$

where  $dr, dx$  are fixed Haar measures on  $S_t^2(W)_{\mathbf{A}}, (W \otimes_F V)_{\mathbf{A}}$ , respectively. Moreover, the  $\mu_r$ 's are tempered measures and for  $\phi \in \mathcal{S}((W \otimes V)_{\mathbf{A}})$ ,  $F_\phi$  is continuous, is an element of  $L^1(S_t^2(W)_{\mathbf{A}})$ , and is the Fourier transform of the function  $F_\phi^*(\cdot)$  on  $S_t^2(W^*)_{\mathbf{A}}$  given by

$$F_\phi^*(\rho) = \int_{(W \otimes V)_{\mathbf{A}}} \phi(x) \psi\left(\frac{1}{2}(T_W(x), \rho)\right) dx.$$

Finally,

$$\text{Et}_W(\phi) = \sum_{r \in S_t^2(W)} \int_{(W \otimes V)_{\mathbf{A}}} \phi d\mu_r, \quad (3.17)$$

the series on the right being absolutely convergent.

Since the convergence of the right hand side of (3.17) is uniform as  $\phi$  varies on a compact subset of  $\mathcal{S}((W \otimes V)_{\mathbf{A}})$ ,  $\text{Et}_W$  is a tempered measure on  $(W \otimes V)_{\mathbf{A}}$ . The formula (3.17) can be restated as:

**Corollary 3.5** We have the identity of the tempered distributions:

$$\text{Et}_W = \sum_{r \in S_t^2(W)} \mu_r. \quad (3.18)$$

## 4 An extension of Weil's abstract lemma

In this section we study the measure  $d\mu_r$  in Theorem 3.4. We begin with a statement of Proposition 1 in [7] (page 6)

**Lemma 4.1** Let  $X$  and  $G$  be two locally compact, abelian groups with fixed Haar measures  $dx, dg$  respectively. Let

$$f : X \rightarrow G$$

be a continuous map such that

(A) For any  $\Phi \in \mathcal{S}(X)$ , the function  $F_\Phi^*$  on  $G^*$  defined by

$$F_\Phi^*(g^*) = \int_X \Phi(x) \{f(x), g^*\} dx, \quad (4.1)$$

(where  $dx \{, \}$  denotes the pairing between  $G$  and  $G^*$ ) is integrable on  $G^*$  and the integral  $\int |F_\Phi^*| dg^*$  (where  $dg^*$  is Haar measure on  $G^*$ ) dual to  $dg$ ) converges uniformly on every compact subset of  $\mathcal{S}(X)$ .

Then one can find a uniquely determined family of positive measures  $\{\mu_g\}_{g \in G}$ , on  $X$ , where  $\text{support}(\mu_g) \subseteq f^{-1}(\{g\})$ , and so that for every continuous function with compact support  $\Phi$  on  $X$ , the function  $F_\Phi$  on  $G$  defined by

$$F_\Phi(g) = \int \Phi d\mu_g, \quad (4.2)$$

is continuous and satisfies

$$\int F_\Phi dg = \int \Phi dx. \quad (4.3)$$

Moreover, the measures  $\mu_g$  are tempered measures and for  $\Phi \in \mathcal{S}(X)$ ,  $F_\Phi$  is continuous, belongs to  $L^1(G)$ , satisfies (4.3), and is the Fourier transform of  $F_\Phi^*$ .

We call  $(X, G, f)$  as in Lemma 4.1, an *admissible triple*.

Let  $M, M_-, M_+, V$  be as in Section 3. And as Section 3,  $W$  denotes a  $F[[t]]$ -submodule of  $M_-$ . For a fixed place  $v$  of  $F$ , we set

$$X_v = (W \otimes V)_{F_v}, \quad G_v = S_t^2(W)_{F_v},$$

and let  $T_v : X_v \rightarrow G_v$  be the  $F_v$ -linear extension of  $T_W$  defined in (3.13). The dual group  $G_v^*$  of  $G_v$  is identified with  $(F_W)_{F_v}$ .

**Lemma 4.2** *If  $\dim V > 6n + 2$ , then above triple  $(X_v, G_v, T_v)$  is an admissible triple, equivalently,*

$$F_\Phi^*(g^*) = \int_X \Phi(x) \{f(x), g^*\} dx,$$

*satisfies condition (A) in Lemma 4.1.*

This lemma is an analog of Proposition 5 in [7] (page 45). We expect that the condition  $\dim V > 6n + 2$  can be replaced by the weaker condition  $\dim V > 6n + 1$ . For our purpose, the condition in the lemma is enough.

*Proof.* For simplicity, we write  $X, G, T$  for  $X_v, G_v, T_v$ . Let  $X_{\mathbf{A}} = (W \otimes V)_{\mathbf{A}}$ ,  $G_{\mathbf{A}} = S_t^2(W)_{\mathbf{A}}$ , and  $T_{\mathbf{A}} : X_{\mathbf{A}} \rightarrow G_{\mathbf{A}}$  be  $T_W \otimes \mathbf{A}$  then we have

$$X_{\mathbf{A}} = X \times X^c, \quad G_{\mathbf{A}} = G \times G^c$$

where  $X^c$  is the restricted product of  $(W \otimes V)_{F_w}$ 's for  $w \neq v$ ,  $G^c$  is the restricted product of  $S_t^2(W)_{F_w}$ 's for  $w \neq v$ . And  $T_{\mathbf{A}} = T \times T_c$ , where  $T_c : X^c \rightarrow G^c$  is defined similarly as  $T$ . Let  $C$  be a compact subset of  $\mathcal{S}(X)$ . We choose a function  $\phi_0 \in \mathcal{S}(X^c)$  such that

$$\int_{X^c} \phi_0(x_c) dx_c \neq 0.$$

Then  $F_{\phi_0}^*(g_c^*)$  given by

$$F_{\phi_0}^*(g_c^*) = \int_{X^c} \phi_0(x_c) \{T_c(x_c), g_c^*\} dx_c,$$

satisfies that  $F_{\phi_0}^*(0) \neq 0$ . Since  $F_{\phi_0}^*(g_c^*)$  is continuous, we have

$$\int_{G^c} |F_{\phi_0}(g_c^*)| dg_c^* = M \neq 0.$$

For each function  $\phi(x) \in \mathcal{S}(X)$ ,  $\phi(x)\phi_0(x_c)$  is in  $\mathcal{S}(X_{\mathbf{A}})$ . Since  $C$  is a compact subset of  $\mathcal{S}(X)$ ,  $C\phi_0$  is a compact subset of  $\mathcal{S}(X_{\mathbf{A}})$ . By Theorem 3.3, and the Proposition 2 in [7], we know that  $X_{\mathbf{A}}, G_{\mathbf{A}}, T_{\mathbf{A}}$  is an admissible triple. We consider the function

$$\int_X \int_{X^c} \phi(x)\phi_0(x_c) \{T(x), g^*\} \{T_c(x_c), g_c^*\} dx dx_c = F_{\phi}(g^*) F_{\phi_0}(g_c^*).$$

Since  $C\phi_0$  is compact, the integral

$$\int_{G^* \times G_c^*} |F_{\phi}(g^*) F_{\phi_0}(g_c^*)| dg^* dg_c^*$$

converges uniformly as  $\phi$  varies over  $C$ . By the Fubini theorem, we have

$$\begin{aligned} & \int_{G^* \times G_c^*} |F_{\phi}(g^*) F_{\phi_0}(g_c^*)| dg^* dg_c^* \\ &= \int_{G^*} |F_{\phi}(g^*)| dg^* \int_{G_c^*} |F_{\phi_0}(g_c^*)| dg_c^* \\ &= M \int_{G^*} |F_{\phi}(g^*)| dg^*. \end{aligned}$$

This implies that

$$\int_{G^*} |F_\phi(g^*)| dg^*$$

converges uniformly as  $\phi$  varies on  $C$ .  $\square$ .

Since  $W$  is a finite dimensional over  $F$  and  $t^N W = 0$  for  $N$  large,  $W$  is isomorphic to

$$F[t]/(t^{k_1})e_1 \oplus \cdots \oplus F[t]/(t^{k_m})e_m$$

as a  $F[[t]]$ -module, where  $e_1, \dots, e_m$  is a quasi-basis of  $W$ . Let  $\bar{W}$  denote the quotient  $W/tW$ , we have

$$\bar{W} \cong F\bar{e}_1 \oplus \cdots \oplus F\bar{e}_m,$$

where  $\bar{e}_i$  is the projection of  $e_i$ . Let  $\bar{G}_v$  denote  $S^2(\bar{W}_{F_v})$ , where  $S^2(\bar{W}_{F_v})$  is the subspace of the symmetric tensors in  $\bar{W}_{F_v} \otimes \bar{W}_{F_v}$ . For simplicity, we shall write  $G, X, \bar{G}$  for  $G_v, X_v, \bar{G}_v$ . We have  $\bar{T}$  given by

$$\bar{T} : \bar{X} \stackrel{\text{def}}{=} (\bar{W} \otimes V)_v \rightarrow \bar{G} \stackrel{\text{def}}{=} S^2(W_v), \quad \sum_i u_i \otimes v_i \mapsto \sum_{i,j} (v_i, v_j) u_i \otimes u_j.$$

The condition  $\dim V > 6n + 2$  implies in particular  $\dim V > 6n + 2 \geq 6m + 2$ ; this implies that the condition for Proposition 5 in [7] (page 45) is satisfied, so  $(\bar{X}, \bar{G}, \bar{T})$  is an admissible triple. The canonical map  $W \rightarrow \bar{W}$  induces surjective linear maps

$$\pi_X : X \rightarrow \bar{X}, \quad \pi_G : G \rightarrow \bar{G}.$$

We have the commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{\pi_X} & \bar{X} \\ \downarrow T & & \downarrow \bar{T} \\ G & \xrightarrow{\pi_G} & \bar{G} \end{array} \quad (4.4)$$

Let  $f : X \rightarrow \bar{G}$  denote  $\pi_G \circ T = \bar{T} \circ \pi_X$ .

**Lemma 4.3** *For  $x \in X$ , the following conditions are equivalent*

- (1).  *$T$  is submersive at  $x$ .*
- (2).  *$f$  is submersive at  $x$ .*
- (3).  *$\bar{T}$  is submersive at  $\pi_X(x)$ .*

*Proof.* Since  $\pi_X$  is linear and surjective, it is submersive at every point. It follows that (2) and (3) are equivalent. Since  $\pi_G$  is linear and surjective, it is submersive at every point, it follows that (1) implies (3). The fact that (3) implies (1) follows directly from Lemma 5.7 in Section 5.

**Lemma 4.4** *The map  $f : X \rightarrow \bar{G}$  satisfies the condition (A) in Lemma 4.1, so  $(X, \bar{G}, f)$  is an admissible triple.*

*Proof.* We use the following diagram to prove the lemma:

$$\begin{array}{ccc} X & \xrightarrow{\pi_X} & \bar{X} \\ & f \searrow & \downarrow \bar{T} \\ & & \bar{G} \end{array}$$

Let  $K$  denote the kernel of  $\pi_X$ . We have a map from  $\mathcal{S}(X) \rightarrow \mathcal{S}(\bar{X})$  given by

$$\Phi \mapsto \bar{\Phi}(\bar{x}) = \int_K \Phi(k + \bar{x}) dk, \quad (4.5)$$

where  $k$  denotes the Haar measure on  $K$ . It is clear that this map is continuous. Consider

$$F_{\Phi}^*(\bar{g}^*) = \int_X \Phi(x) \langle f(x), \bar{g}^* \rangle dx$$

In the right hand side, we integrate over  $K$  first, and notice that

$$\langle f(x), \bar{g}^* \rangle = \langle \bar{T}(\bar{x}), \bar{g}^* \rangle$$

where  $\bar{x} = \pi_X(x)$ , we get

$$F_{\Phi}^*(\bar{g}^*) = \int_{\bar{X}} \bar{\Phi}(\bar{x}) \langle \bar{T}(\bar{x}), \bar{g}^* \rangle d\bar{x},$$

since  $(\bar{X}, \bar{G}, \bar{T})$  is an admissible triple, the right hand side is in  $L^1(\bar{G})$ . And if  $\Phi$  runs through a compact subset of  $\mathcal{S}(X)$ , then  $\bar{\Phi}$  which is related to  $\Phi$  by (4.5) runs through a corresponding compact subset of  $\mathcal{S}(\bar{X})$ , so the integral

$$\int |F_{\Phi}^*(\bar{g}^*)| d\bar{g}^*$$

converges uniformly. This proves the lemma.  $\square$



By Lemma 4.1, we have a family of measures  $\{\mu_{\bar{g}}\}_{\bar{g} \in \bar{G}}$  on  $X$ , with  $\text{support}(\mu_{\bar{g}}) \subset f^{-1}(\bar{g})$ , such that for every  $\Phi \in C_c(X)$ ,  $\int \Phi d\mu_{\bar{g}}$  is continuous function of  $\bar{g}$  and

$$\int_{\bar{G}} \left( \int \Phi d\mu_{\bar{g}} \right) d\bar{g} = \int_X \Phi dx. \quad (4.6)$$

On the other hand, apply Lemma 4.1 to the admissible triple  $(\bar{G}, \bar{X}, \bar{T})$ , we have a family of measures  $\{\mu_{\bar{g}}^{\bar{T}}\}_{\bar{g} \in \bar{G}}$  on  $\bar{X}$ , with  $\text{support}(\mu_{\bar{g}}^{\bar{T}}) \subset \bar{T}^{-1}(\bar{g})$ , and

$$\int_{\bar{G}} \left( \int \bar{\Phi} d\mu_{\bar{g}}^{\bar{T}} \right) d\bar{g} = \int_{\bar{X}} \bar{\Phi} d\bar{x}. \quad (4.7)$$

Suppose that  $\Phi$  and  $\bar{\Phi}$  are related by (4.5) and the Haar measures  $dg, d\bar{g}, dk$  are compatible so that the right hand sides of (4.6) and (4.7) are equal. We then have

$$\int_{\bar{G}} \left( \int \Phi d\mu_{\bar{g}} \right) d\bar{g} = \int_{\bar{G}} \left( \int \bar{\Phi} d\mu_{\bar{g}}^{\bar{T}} \right) d\bar{g} \quad (4.8)$$

We claim that the truth of (4.8) for all  $\Phi \in C_c(X)$  implies that

$$\int \Phi d\mu_{\bar{g}} = \int \bar{\Phi} d\mu_{\bar{g}}^{\bar{T}}. \quad (4.9)$$

To prove this, take arbitrary  $h(\bar{g}) \in C_c(\bar{G})$ , let  $f^*h = h \circ f$ , replace  $\Phi$  in (4.8) by  $f^*h\Phi$ . We get

$$\int_{\bar{G}} h(\bar{g}) \left( \int \Phi d\mu_{\bar{g}} \right) d\bar{g} = \int_{\bar{G}} h(\bar{g}) \left( \int \bar{\Phi} d\mu_{\bar{g}}^{\bar{T}} \right) d\bar{g}.$$

The above is true for all  $h(\bar{g}) \in C_c(\bar{G})$ , and  $\int \Phi d\mu_{\bar{g}}, \int \bar{\Phi} d\mu_{\bar{g}}^{\bar{T}}$  are continuous functions of  $\bar{g}$ , so we have (4.9). We rewrite (4.9) as

$$\int \Phi d\mu_{\bar{g}} = \int \left( \int_K \Phi(k + \bar{x}) dk \right) d\mu_{\bar{g}}^{\bar{T}} \quad (4.10)$$

Recall Lemma 17 [7] (page 52), the support of  $\mu_{\bar{g}}^{\bar{T}}$  is on the  $\bar{T}^{-1}(\bar{g})_{\text{re}}$  (the regular points (= submersive points) in  $\bar{T}^{-1}(\bar{g})$ ). By (4.10), the support of  $\mu_{\bar{g}}$  is in  $\pi_X^{-1}\bar{T}^{-1}(\bar{g})_{\text{re}}$ , which is precisely the set of the regular points in  $f^{-1}(\bar{g})$  by Lemma 4.3. We have proved

**Lemma 4.5** *The measure  $\mu_{\bar{g}}$  is supported on  $f^{-1}(\bar{g})_{\text{re}}$ , the subset of regular points of  $f^{-1}(\bar{g})$ .*

We consider the diagram

$$\begin{array}{ccc} X & & \\ \downarrow T & \searrow f & \\ G & \xrightarrow{\pi_G} & \bar{G} \end{array}$$

For the admissible triple  $(X, G, T)$ , Lemma 4.1 implies that we have a family of measures  $\mu_g$  ( $g \in G$ ) supported on  $T^{-1}(g)$  such that for  $\Phi \in C_c(X)$ ,  $F_\Phi(g) \stackrel{\text{def}}{=} \int \Phi d\mu_g \in C(G)$ , we have

$$\int_G F_\Phi(g) dg = \int_X \Phi(x) dx \quad (4.11)$$

We take a subspace of  $G$  that maps isomorphically onto  $\bar{G}$ , we denote this space by  $\bar{G}$ , so we have the identification  $G = K \times \bar{G}$ , where  $K$  is the kernel of  $\pi_G$ . Since  $\Phi$  has compact support,  $F_\Phi$  has compact support, and it is continuous, so we have

$$\bar{g} \rightarrow \int_K F_\Phi(\bar{g} + k) dk$$

is in  $C(\bar{G})$ . The left hand side of (4.11) can be written as

$$\int_{\bar{G}} \int_K F_\Phi(\bar{g} + k) dk d\bar{g},$$

so we have

$$\int_{\bar{G}} \int_K F_\Phi(\bar{g} + k) dk d\bar{g} = \int_X \Phi(x) dx \quad (4.12)$$

On the other hand, use the triple  $(X, \bar{G}, f)$ , we have by (4.6

$$\int_{\bar{G}} \left( \int \Phi d\mu_{\bar{g}} \right) d\bar{g} = \int_X \Phi(x) dx \quad (4.13)$$

Comparing (4.12) and (4.13), we get

$$\int_{\bar{G}} \int_K F_\Phi(\bar{g} + k) dk d\bar{g} = \int_{\bar{G}} \left( \int \Phi d\mu_{\bar{g}} \right) d\bar{g}. \quad (4.14)$$

Take an arbitrary  $h \in C_c(\bar{G})$ , let  $f^*h = h \circ f \in C(X)$ , and replacing  $\Phi$  in (4.14) by  $f^*h\Phi$ , we get

$$\int_{\bar{G}} h(\bar{g}) \int_K F_\Phi(\bar{g} + k) dk d\bar{g} = \int_{\bar{G}} h(\bar{g}) \left( \int \Phi d\mu_{\bar{g}} \right) d\bar{g}.$$

This is true for arbitrary  $h \in C_c(\bar{G})$ , and both  $\int_K F_\Phi(\bar{g} + k)dk$  and  $\int \Phi d\mu_{\bar{g}}$  are continuous functions, so we have

$$\int_K F_\Phi(\bar{g} + k)dk = \int \Phi d\mu_{\bar{g}}.$$

But the measure  $\mu_{\bar{g}}$  is supported on  $f^{-1}(\bar{g})_{\text{re}}$ , so it is a gauge measure (see section 5 of [7] for the definition of "gauge" measure),

$$f^{-1}(\bar{g}) = T^{-1}(\pi_G^{-1}(\bar{g})) = T_X^{-1}(g + K)$$

Note that

$$f^{-1}(\bar{g})_{\text{re}} = \cup_{k \in K} T^{-1}(g + k)_{\text{re}}$$

For each given  $\bar{g}$ ,  $T^{-1}(\bar{g} + k)_{\text{re}}$  is non-singular subvariety of  $X$ , we have a gauge form  $d\delta_k$  on it, we have

$$\int \Phi d\mu_{\bar{g}} = \int_K \int_{T^{-1}(\bar{g}+k)} \Phi d\delta_k dk$$

we obtain

$$\int_K F_\Phi(\bar{g} + k)dk = \int_K \int T^{-1}(\bar{g} + k) \Phi(x) d\delta_k dk. \quad (4.15)$$

The above holds for arbitrary  $\Phi \in C_c(X)$ , use the same method we used to deduce (4.9) from (4.8), we deduce from (4.15) that

$$F_\Phi(\bar{g} + k) = \int T^{-1}(\bar{g} + k)_{\text{re}} \Phi d\delta_k$$

So we have proved

**Lemma 4.6** *The measure  $\mu_{g,v}$  for the triple  $(X_v, G_v, T_v)$  is supported on  $T_v^{-1}(g)_{\text{re}}$  and is the gauge measure.*

Lets recall the meaning of "gauge" measure ([7], section 5). In the situation as Lemma 4.6. We first take an invariant top form  $\eta$  on  $G$  and an invariant top form  $\omega$  on  $X$ . Let  $X'$  be the open set of  $X$  that consists of all the points where  $T$  is submersive. Near each point  $x \in X'$ , there is a form  $\theta_x$  such that  $\theta \wedge T^* \eta = \omega$ . For each  $y \in G$ , the local forms  $\theta$ , restrict to  $T^{-1}(y)_{\text{re}} = T^{-1}(y) \cap X'$ , give a top form  $\theta_y$  on  $T^{-1}(y)_{\text{re}}$ , which defines a measure which is equal to  $\mu$  in Lemma 4.6.

Now we consider the global situation. Let  $X, G, T_W$  as in (3.16). For each  $r \in S_t^2(W)$ , we consider the inverse image  $T_W^{-1}(r)$ . Notice that  $T$  is submersive at a generic point, the space  $X'$  formed by the points at which  $f$  is submersive is  $F$ -open in  $X$ . We take a top form  $\eta$  over  $G$  and a top form  $\omega$  on  $X$ , we assume that the Tamagawa measures on  $G(\mathbf{A})$  ( $X_{\mathbf{A}}$  resp.) with respect to  $\eta$  (resp  $\omega$ ) are the Haar measure normalized by the condition that the covolume of  $G(F)$  ( $X(F)$ ) is 1. The space  $X'$  can be covered by  $F$ -open subsets  $U_\lambda$  such that, there is a form  $\theta_\lambda$  rational over  $F$  satisfying  $\theta \wedge f^* = \omega$ . For each  $i \in G(F)$ , then the  $\theta$ 's restrict on  $f^{-1}(i) \cap X'$  to get a top form  $\theta_i$  on  $f^{-1}(i) \cap X'$ . By Lemma 4.6,  $d\mu_{i,v}$  is given by  $|\theta_i|_v$ . Using a similar argument as in [7], Section 42, we can prove that 1 is a system of convergence factor of  $|\theta_i|_{\mathbf{A}}$ . And we have

**Theorem 4.7** *For each  $r \in S_t^2(W)$ , the measure  $\mu_r$  in (3.17) is supported in  $T_W^{-1}(r)_{\text{re}}$  and it is the same as  $|\theta_r|_{\mathbf{A}}$ , the measure define by the form  $\theta_r$ .*

## 5 Classification of orbits of orthogonal groups

As in Section 3, we denote  $M$  an snt-module with decomposition  $M = M_- \oplus M_+$  into  $t$ -Lagrangian subspaces and  $V$  a finite dimensional vector space over  $F$  with a non-degenerate, bilinear symmetric form  $(,)$ . The orthogonal group  $G(F[[t]])$  acts on  $M \otimes V$ , leaving the subspace  $M_- \otimes V$  invariant. The purpose of this section to give a complete set of invariants of  $G(F[[t]])$ -orbits in  $M_- \otimes V$  (see Theorem 5.4 below).

**Definition 5.1** *Let  $W$  be a finitely generated  $F[[t]]$ -module. A submodule  $L \subset W$  is called a primitive submodule if one of the following equivalent conditions is satisfied:*

- (1). *there is a complement  $F[[t]]$ -submodule  $L'$ , i.e.  $W = L \oplus L'$ .*
- (2). *the natural map  $L/tL \rightarrow W/tW$  induced from the embedding  $L \hookrightarrow M$  is injective.*

*Examples.* (1) Let  $W = F[[t]]/(t^{k_1}) \oplus \cdots \oplus F[[t]]/(t^{k_m})$ , For any  $l \leq m$ ,  $L = F[[t]]/(t^{k_1}) \oplus \cdots \oplus F[[t]]/(t^{k_l})$  is a primitive submodule of  $M$ . (2). Let  $F[[t]]^m$  be a free  $F[[t]]$ -module of rank  $m$ , then  $F[[t]]^l$  (consists of elements with last  $(m-l)$ -components 0) is a primitive submodule. (3).  $\{0\}$  is a primitive submodule for any  $M$ .

Let  $W$  be a finitely generated  $F[[t]]$ -module,  $e_1, \dots, e_m$  is called a *quasi-basis* of  $W$  if every  $e_i \neq 0$ , every element  $x \in W$  can be written as a  $F[[t]]$ -linear combination of  $e_1, \dots, e_m$ , and  $a_1 e_1 + \dots + a_m e_m = 0$  ( $a_i \in F[[t]]$ ) implies that all  $a_i e_i = 0$ . If  $W$  is a finite dimensional  $F[[t]]$ -module, the "quasi-basis" defined above is the same notion as defined in Section 2. If  $W$  is a free  $F[[t]]$ -module, then a quasi-basis of  $W$  is the same as a basis of  $W$ . In the example (1) above, there is a quasi-basis of  $m$  elements. It is clear that  $e_1, \dots, e_m$  is a quasi-basis of  $W$  if and only if the images of  $e_1, \dots, e_m$  in  $W/tW$  form an  $F$ -basis of vector space  $W/tW$ . Therefore any two quasi-bases have the same number of elements. The cardinality of a quasi-basis is called the *rank* of  $W$ .

Every

$$x = \sum_i u_i \otimes v_i \in M_- \otimes V = M_- \otimes_{F[[t]]} V[[t]]$$

gives rise to an  $F[[t]]$ -linear map

$$f_x : V[[t]] \rightarrow M_-, \quad f_x(v) = \sum_i (v_i, v) u_i. \quad (5.1)$$

We denote by  $\text{Im } f_x$  the image of  $f_x$ , which is an  $F[[t]]$ -submodule of  $M_-$ . Let  $e_1, \dots, e_m$  be a quasi-basis of  $\text{Im } f_x$ , then

$$\text{Im } f_x \cong F[[t]]/(t^{k_1})e_1 \oplus \dots \oplus F[[t]]/(t^{k_m})e_m, \quad (5.2)$$

where  $k_i$  is the smallest positive integer such that  $t^{k_i} e_i = 0$ .

**Lemma 5.2** *Let  $e_1, \dots, e_m$  be a quasi-basis of  $\text{Im } f_x$ , and suppose  $v_1, \dots, v_m \in V[[t]]$  satisfy  $f_x(v_i) = e_i$  ( $i = 1, \dots, m$ ), then  $\text{Span}_{F[[t]]}\{v_1, \dots, v_m\}$  is a primitive submodule of  $V[[t]]$  and  $v_1, \dots, v_m$  is a basis of  $\text{Span}_{F[[t]]}\{v_1, \dots, v_m\}$ .*

*Proof.* Set  $L = \text{Span}_{F[[t]]}\{v_1, \dots, v_m\}$ , then  $f_x|_L : L \rightarrow \text{Im } f_x$  induces a linear isomorphism:

$$\bar{f}_x : L/tL \rightarrow \text{Im } f_x/t(\text{Im } f_x).$$

This implies in particular,  $\dim L/tL = m$ , so the map  $L/tL \rightarrow V[[t]]/tV[[t]]$  induced from  $L \subset V[[t]]$  is injective, so  $L$  is primitive submodule of  $V[[t]]$ . The other conclusions are clear.  $\square$

The bilinear form  $(,)$  on  $V$  can be extended to a  $F[[t]]$ -valued bilinear form on  $V[[t]] = V \otimes_F F[[t]]$ . This  $F[[t]]$ -valued bilinear form on  $V[[t]]$  is non-degenerate. It is easy to prove

**Lemma 5.3** *Let  $e_1, \dots, e_m$  be a quasi-basis of  $\text{Im } f_x$ , then there are elements  $w_1, \dots, w_m \in V[[t]]$  such that*

*(1)  $\text{Span}_{F[[t]]}\{w_1, \dots, w_m\}$  is a primitive submodule of  $V[[t]]$  and  $w_1, \dots, w_m$  is basis of  $\text{Span}_{F[[t]]}\{w_1, \dots, w_m\}$ .*

*(2)*

$$x = e_1 \otimes w_1 + \dots + e_m \otimes w_m.$$

*Proof.* Choose  $v_i \in V[[t]]$  ( $i = 1, \dots, m$ ) such that  $f_x(v_i) = e_i$ . By Lemma 5.2,  $\text{Span}_{F[[t]]}\{v_1, \dots, v_m\}$  is a primitive submodule of  $V[[t]]$  and  $v_1, \dots, v_m$  is a basis. It is clear that

$$V[[t]] = \text{Span}_{F[[t]]}\{v_1, \dots, v_m\} \oplus \ker(f_x).$$

Let  $v_{m+1}, \dots, v_N$  be a basis of  $\ker(f_x)$ . Then  $v_1, \dots, v_N$  is basis of  $V[[t]]$ . Now let  $w_1, \dots, w_N$  be the dual basis of  $v_1, \dots, v_N$ , i.e.,  $(v_i, w_j) = \delta_{ij}$ . This is clear that  $v = \sum_{i=1}^m e_i \otimes w_i$ .  $\square$

From Lemma 5.3, we see that  $x \in \text{Im } f_x \otimes_{F[[t]]} V[[t]]$ . We define a map

$$T : \text{Im } f_x \otimes_{F[[t]]} V[[t]] \rightarrow S_t^2(\text{Im } f_x), \quad \sum_i^N u_i \otimes v_i \mapsto \sum_{i,j=1}^N (v_i, v_j) u_i \otimes u_j.$$

Where  $S_t^2(\text{Im } f_x)$  denote the subspace of symmetric tensors in  $\text{Im } f_x \otimes_{F[[t]]} \text{Im } f_x$ . We remark that though  $\text{Im } f_x$  is a submodule of  $M_-$ , but the natural map  $S_t^2(\text{Im } f_x) \rightarrow S_t^2(M_-)$  is in general *not* an embedding.

**Theorem 5.4** *Two elements  $x, y \in M_- \otimes V$  are in the same  $G(F[[t]])$ -orbit iff  $\text{Im } f_x = \text{Im } f_y$  and  $T(x) = T(y)$ .*

We need some preparations for proving the theorem.

We recall a special case of Witt's theorem (see [4]):

**Theorem 5.5** *If  $L_1, L_2 \subseteq V[[t]]$  are two primitive submodules of  $V[[t]]$  and if  $\sigma : L_1 \rightarrow L_2$  is an isometry, then  $\sigma$  can be extended to an isometry in  $G(F[[t]])$ .*

It is clear that if  $x, y$  are in the same  $G(F[[t]])$ -orbit, then  $\text{Im } f_x = \text{Im } f_y$  and  $T(x) = T(y)$ . Conversely, if  $\text{Im } f_x = \text{Im } f_y \stackrel{\text{def}}{=} W$  and  $T(x) = T(y)$ . Let  $e_1, \dots, e_m$  be a quasi-basis for  $W$ , and  $W$  be as (5.2). We may assume that

$k_1 \geq k_2 \geq \cdots \geq k_m$ . Then  $S_t^2(W)$  has a quasi-basis  $e_{ij} \stackrel{\text{def}}{=} e_i \otimes e_j + e_j \otimes e_i$  ( $1 \leq i \leq j \leq m$ ), and

$$S_t^2(W) = \sum_{1 \leq i \leq j \leq m} F[[t]]/(t^{k_j})e_{ij}.$$

By lemma 5.3, we may write

$$x = e_1 \otimes a_1 + \cdots + e_m \otimes a_m, \quad y = e_1 \otimes b_1 + \cdots + e_m \otimes b_m$$

with  $\{a_1, \dots, a_m\}$  and  $\{b_1, \dots, b_m\}$  satisfy condition (1) in Lemma 5.3. Then  $T(x) = T(y)$  implies that

$$(a_i, a_j) = (b_i, b_j) \pmod{t^{\min(k_i, k_j)}}. \quad (5.3)$$

**Lemma 5.6** *Let  $L_1$  and  $L_2$  be two primitive submodules of  $V[[t]]$  with bases  $a_1, \dots, a_m$  and  $b_1, \dots, b_m$ . Let  $1 \geq k_1 \geq \cdots \geq k_m$ . If (5.3) holds, then the set  $b_1, \dots, b_m$  can be altered to another set  $\tilde{b}_1, \dots, \tilde{b}_m$  such that*

$$\tilde{b}_i = b_i \pmod{t^{k_i}}, \quad \text{for } 1 \leq i \leq m, \quad (5.4)$$

and

$$(\tilde{b}_i, \tilde{b}_j) = (a_i, a_j) \quad \text{for } 1 \leq i, j \leq m, \quad (5.5)$$

and the  $F[[t]]$ -span of  $\tilde{b}_1, \dots, \tilde{b}_m$  is a primitive submodule of  $V[[t]]$  with  $\tilde{b}_1, \dots, \tilde{b}_m$  as a basis.

Suppose the truth of Lemma 5.6, then Theorem 5.4 can be proved as follows. The equation (5.4) implies that

$$y = \sum e_i \otimes b_i = \sum e_i \otimes \tilde{b}_i.$$

The equation (5.5) implies that the map  $\sigma : L_1 \rightarrow L_2$  give by  $a_i \mapsto \tilde{b}_i$  is an isometry, by Theorem 5.5,  $\sigma$  can be extended to  $g \in G(F[[t]])$ . then

$$y \cdot g = (\sum e_i \otimes \tilde{b}_i) \cdot g = \sum e_i \otimes \sigma(\tilde{b}_i) = \sum e_i \otimes a_i = x.$$

It remains to prove Lemma 5.6.

*Proof of Lemma 5.6.* We use induction on  $m$ . For case  $m = 1$ , we first take  $c \in V[[t]]$  such that  $(b_1, c) = 1$ , we want to find  $\tilde{b}_1 = b_1 + t^{k_1}h(t)c$  where  $h(t) = h_0 + h_1t + h_2t^2 + \cdots \in F[[t]]$  such that We have

$$(a_1, a_1) = (\tilde{b}_1, \tilde{b}_1)$$

which is equivalent to

$$(a_1, a_1) - (b_1, b_1) = 2t^{k_1}h(t) + t^{2k_1}h(t)^2(c, c). \quad (5.6)$$

Since  $(a_1, a_1) = (b_1, b_1) \bmod t^{k_1}$ , we see that 5.6 holds  $\bmod t^{k_1}$  for arbitrary  $h(t)$ . Compare the coefficient of  $t^{k_1}$ , we solve for  $h_0$ , after  $h_0$ , we compare coefficient of  $t^{k_1+1}$ , we solve  $h_1$ . It is clear that the similar process can be continued to solve all  $h_i$ . Assume the Lemma holds for  $m-1$ , so we can find  $\tilde{b}_1, \dots, \tilde{b}_{m-1}$  such that

$$\tilde{b}_i \cong b_i \bmod t^{k_i}, \text{ for } 1 \leq i \leq m-1, \quad (5.7)$$

and

$$(\tilde{b}_i, \tilde{b}_i) = (a_i, a_i) \text{ for } 1 \leq i, j \leq m-1. \quad (5.8)$$

We may assume  $b_i = \tilde{b}_i$  for  $i = 1, \dots, m-1$ . Since  $(\ )$  on  $V[[t]]$  is non-degenerate, we can find  $c_1, c_2, \dots, c_m \in V[[t]]$  such that

$$(b_i, c_j) = \delta_{i,j}. \quad (5.9)$$

We want to find  $h_1(t), \dots, h_m(t) \in F[[t]]$  such that

$$\tilde{b}_m = b_m + t^{k_m}(h_1(t)c_1 + \dots + h_m(t)c_m)$$

satisfies the following  $m$  equations

$$(\tilde{b}_i, \tilde{b}_m) = (a_i, a_m), \quad i = 1, \dots, m \quad (5.10)$$

Let  $h_i(t) = \sum_{s=0}^{\infty} h_{i,s}t^s$ . The equations (5.10) already hold  $\bmod t^{k_m}$ . Compare the coefficient of  $t^{k_m}$  of (5.10), we get a linear system with  $m$ -variables  $h_{1,0}, \dots, h_{m,0}$  and  $m$  equations, this system has non-zero determinant, thanks to (5.9), we can solve for  $h_{1,0}, \dots, h_{m,0}$ . Then we compare coefficient of  $t^{k_m+1}$  of (5.10), we get a linear system with  $m$  equations and  $m$  variables  $h_{1,1}, \dots, h_{m,1}$ , and again because of (5.9), the system has a solution. It is clear that this process can be continued to solve for all  $h_{i,s}$ 's.  $\square$

**Lemma 5.7** *Let  $v$  be a place of  $F$ ,  $W$  be a  $F[[t]]$ -submodule of  $M_-$ , let  $W_v = W \otimes F_v, V_v = V \otimes F_v$ . The map  $T : W_v \otimes V_v \rightarrow S_t^2(W_v)$  given by*

$$T\left(\sum_i u_i \otimes v_i\right) = \sum_{i,j} (v_i, v_j) u_i \otimes u_j$$

*is submersive at  $x_0 \in W_v \otimes V_v$  iff  $\text{Im } f_{x_0} = W_v$*



*Proof.* For simplicity, we denote  $W_v, V_v, F_v$  by  $W, V, F$  respectively. Let  $a_1, \dots, a_m$  be a quasi-basis of the  $F[[t]]$ -module  $\text{Im } f_{x_0}$ . By Lemma 5.3,  $x_0$  can be written as

$$x_0 = \sum_i a_i \otimes b_i = a_1 \otimes b_1 + \dots + a_m \otimes b_m$$

where  $b_1, \dots, b_m$  is a basis of  $\text{Span}\{b_1, \dots, b_m\}$  and  $\text{Span}\{b_1, \dots, b_m\}$  is a primitive  $F[[t]]$ -submodule of  $V[[t]]$ . We first find a formula for the tangent map

$$dT_{x_0} : T_{x_0} = W \otimes_{F[[t]]} V[[t]] \rightarrow T_{y_0} = S_t^2(W)$$

where  $y_0 = T(x_0)$ . Take a line  $x(\epsilon) = \sum_i a_i \otimes b_i + \epsilon \sum_j u_j \otimes v_j$  passing through  $x_0$  in the direction  $\sum_j u_j \otimes v_j$ , Then

$$T(x(\epsilon)) = \sum (b_i, b_j) a_i \otimes a_j + \epsilon \sum (b_i, v_j) (a_i \otimes u_j + u_j \otimes a_i) + \epsilon^2 \sum (v_i, v_j) u_i \otimes u_j.$$

So we have

$$dT_{x_0}(\sum_j u_j \otimes v_j) = \sum (b_i, v_j) (a_i \otimes u_j + u_j \otimes a_i). \quad (5.11)$$

From this formula, we see that  $dT_{x_0}$  is  $F[[t]]$ -linear. If  $\text{Im } f_{x_0} \neq W$ , then  $\text{Im } dT_{x_0} \subset \text{Im } f_{x_0} \otimes W + \text{Im } f_{x_0} \otimes W$ ,  $dT_{x_0}$  is not surjective, i.e.  $T$  is not submersive at  $x_0$ . If  $\text{Im } f_{x_0} = W$ , since  $(\cdot)$  on  $V[[t]]$  is non-degenerate, for each  $1 \leq k \leq m$ , we can find  $v \in V[[t]]$  such that  $(b_i, v) = \delta_{i,k}$ , then  $dT_{x_0}(a_l \otimes v) = a_l \otimes a_k + a_k \otimes a_l$ . So all  $a_l \otimes a_k + a_k \otimes a_l$  are in  $\text{Im } f_{x_0}$ . So  $dT_{x_0}$  is surjective and  $T$  is submersive at  $x_0$ .  $\square$

For an  $F[[t]]$ -submodule  $W$  of  $M_-$ ,  $T_W : W \otimes V \rightarrow S_t^2(W)$  as in Section 3. For  $i \in S_t^2(W)$ , We denote by  $U(i)$  the variety of elements in  $T_W^{-1}(i)$  where  $T_W$  is submersive. Of course the set of  $F$ -points  $U(i)_F$  may be empty. By Lemma 5.7, Theorem 5.4 can be reformulated as

**Theorem 5.8** *The  $G(F[[t]])$ -orbits in  $M_- \otimes V$  are in one-to-one correspondence with the set of pairs  $(W, i)$  with  $W \in \text{Gr}(M_-, t)$ ,  $i \in S_t^2(W)$  such that  $U(i)_F$  is not empty. The correspondence is the following, for  $x \in M_- \otimes V$ , its orbit corresponds to the pair  $(W, i)$ , where  $W = \text{Im } f_x$ , and  $i = T_W(x)$ .*

## 6 Theta series.

We continue with the snt-module  $M$  with decomposition  $M = M_- \oplus M_+$  and  $V$  a finite dimensional vector space with a non-degenerate, bilinear symmetric form  $(, )$  as in Section 3. For each  $\phi \in \mathcal{S}(X_{\mathbf{A}}) = \mathcal{S}((M_- \otimes V)_{\mathbf{A}})$ , the theta functional  $\theta(\phi)$  is defined by (3.7). Recall  $G^q(\mathbf{A}[[t]])$  acts on  $\mathcal{S}(X_{\mathbf{A}})$ , the action formula is given as follows. An element  $g \in G^q(\mathbf{A}[[t]])$  has the block decomposition

$$\begin{bmatrix} \alpha_g & \beta_g \\ \gamma_g & \delta_g \end{bmatrix}$$

with respect the decomposition

$$(M \otimes V)_{\mathbf{A}} = (M_- \otimes V)_{\mathbf{A}} \oplus (M_+ \otimes V)_{\mathbf{A}}.$$

Since  $G^q(\mathbf{A}[[t]])$  preserves  $(M_{\pm} \otimes V)_{\mathbf{A}}$ , so we have  $\beta_g = 0$  and  $\gamma_g = 0$ . Then the action of  $g$  on  $\mathcal{S}((M_- \otimes V)_{\mathbf{A}})$  is given by

$$(g \cdot \phi)(x) = \phi(x\alpha_g) = \phi(xg) \quad (6.1)$$

If  $g \in G^q(F[[t]])$ , it is clear that

$$\theta(g \cdot \phi) = \theta(\phi).$$

So  $\theta(g \cdot \phi)$  is a continuous function on  $G^q(F[[t]]) \backslash G^q(\mathbf{A}[[t]])$ . Let  $dg$  be the Haar measure on  $G^q(\mathbf{A}[[t]])$  such that the volume of  $G^q(F[[t]]) \backslash G^q(\mathbf{A}[[t]])$  is 1.

**Lemma 6.1** *If  $(V, (, ))$  is anisotropic over  $F$  or  $\dim V - r > \frac{1}{2}\dim M + 1$ , where  $r$  is the dimension of a maximal isotropic subspace of  $V$ , then the integral*

$$\text{It}(\phi) \stackrel{\text{def}}{=} \int_{G^q(F[[t]]) \backslash G^q(\mathbf{A}[[t]])} \theta(g \cdot \phi) dg \quad (6.2)$$

*converges*

*Proof.* If  $(V, (, ))$  is anisotropic over  $F$ , then  $G^q(F[[t]]) \backslash G^q(\mathbf{A}[[t]])$  is compact, so (6.2) converges. Let  $U$  be the unipotent radical of  $G^q$ , then  $G^q = G \ltimes U$ . Let  $D$  be a compact fundamental domain of  $U(F) \backslash U(\mathbf{A})$ . Then

$$(6.2) = \int_{G(F) \backslash G^q(A)} \int_{a \in D} \sum_{r \in M_- \otimes V} \phi(rag) dadg = \int_{G(F) \backslash G^q(A)} \theta(g \cdot \bar{\phi}) dg, \quad (6.3)$$

where  $\bar{\phi}(x) = \int_D \phi(xa)da$ . Since  $D$  is compact,  $\bar{\phi} \in \mathcal{S}((M_- \otimes V)_{\mathbf{A}})$ . By the convergence criterion in Proposition 8 [7], the right hand side of (6.3) converges under the condition  $\dim V - r > \frac{1}{2}\dim M + 1$ .  $\square$

We can write the integral (6.2) as a sum of orbital integrals. Let  $\mathcal{O}$  be a set of representatives of  $G(F[[t]])$ -orbit in  $M_- \otimes V$ . For each  $\xi \in \mathcal{O}$ , let  $G_\xi$  denote its isotropy subgroup in  $G^q(F[[t]])$ . Then the integral (6.2) can be written as

$$\int_{G^q(F[[t]]) \backslash G^q(\mathbf{A}[[t]])} \sum_{\xi \in \mathcal{O}} \sum_{\tau \in G_\xi \backslash G^q(F[[t]])} \phi(\xi \tau g) dg, \quad (6.4)$$

which can be further written as

$$\sum_{\xi \in \mathcal{O}} \text{vol}(G_\xi \backslash G_{\xi, \mathbf{A}}) \int_{G_{\xi, \mathbf{A}} \backslash G^q(\mathbf{A}[[t]])} \phi(\xi g) dg \quad (6.5)$$

and we have thereby expressed  $\text{It}(\phi)$  as

$$\text{It}(\phi) = \sum_{\xi \in \mathcal{O}} \text{vol}(G_\xi \backslash G_{\xi, \mathbf{A}}) \int_{G_{\xi, \mathbf{A}} \backslash G^q(\mathbf{A}[[t]])} \phi(\xi g) dg. \quad (6.6)$$

## 7 Siegel-Weil formula

We assume in this section  $M$  and  $V$  satisfies the conditions that  $\dim V > 6n + 2$ , where  $n$  is the number of  $H_k$ 's in the decomposition of  $M$  as in (3.4) and  $V$  satisfies the conditions in Lemma 6.1. By Theorem 3.3, the  $t$ -Eisenstein series  $\phi \in \mathcal{S}((M_- \otimes V)_{\mathbf{A}}) \mapsto \text{Et}(\phi)$  is a tempered distribution on  $(M_- \otimes V)_{\mathbf{A}}$ . And for each  $W \in \text{Gr}(M_-, t)$ , we have a tempered distribution  $\phi \mapsto \text{Et}_W(\phi)$ , given by (3.12). We have

$$\text{Et} = \sum_{W \in \text{Gr}(M_-, t)} \text{Et}_W.$$

By Theorem 4.7,

$$\text{Et}_W = \sum_{i \in \text{St}^2(W)} \mu_i.$$

We denote  $\mu_i$  by  $\text{Et}_{W,i}$ . Therefore we have

$$\text{Et} = \sum_{W \in \text{Gr}(M_-, t)} \sum_{i \in \text{St}^2(W)} \mu_{W,i}. \quad (7.1)$$

Moreover the measure  $\text{Et}_{W,i} = \mu_i$  is the gauge measure as described in Theorem 4.7, which implies in particular  $\mu_i$  is 0 if  $U(i)_{\mathbf{A}}$  is empty.

On the other hand,

$$\phi \in \mathcal{S}((M_- \otimes V)_{\mathbf{A}}) \mapsto \text{It}(\phi)$$

given in (6.2) is a tempered distribution and it has a decomposition given by (6.6). By Theorem 5.8, each orbit corresponds uniquely to a pair  $(W, i)$  where  $W \in \text{Gr}(U_-, t)$  and  $i \in \text{St}^2(W)$ . So we may write

$$\text{It} = \sum_{W,i} \text{I}_{W,i}$$

where

$$\text{I}_{W,i}(\phi) = \text{vol}(G_{\xi} \backslash G_{\xi, \mathbf{A}}) \int_{G_{\xi, \mathbf{A}} \backslash G^q(\mathbf{A}[[t]])} \phi(\xi g) dg.$$

where  $(W, i)$  corresponds to the orbit containing  $\xi$ , i.e.,  $\text{Im } f_{\xi} = W$  and  $T_W(\xi) = i$ . If  $U(i)$  is empty,  $(W, i)$  doesn't correspond to any orbit, in this case we define

$$\text{I}_{W,i} \stackrel{\text{def}}{=} 0.$$

We shall prove that  $\text{Et} = \text{It}$ , and actually we shall prove more:  $\text{Et}_{W,i} = \text{It}_{W,i}$  for any pair  $(W, i)$ . We use the induction on  $\dim M$ . The case  $\dim M = 2$  is the classical result in [7].

Our proof is entirely parallel to that of [7]. To start with, we introduce some notations. Let  $\pi : \widehat{Sp}_{2N}(\mathbf{A}) \rightarrow Sp_{2N}(\mathbf{A})$  denote the double cover (recall  $2N = \dim M \dim V$ ). Let  $\widehat{Sp}(M, t)_{\mathbf{A}}$  denote  $\pi^{-1}(Sp(M, t)_{\mathbf{A}})$ . We let

$$M(k) = \oplus_{k_i=k} H_{k_i},$$

so

$$M = M(l_1) \oplus \cdots \oplus M(l_s), \quad l_1 > \cdots > l_s.$$

Recall Corollary 2.6,  $Sp(M, t) = N \ltimes H$ , where  $N$  is the unipotent radical, and

$$H = \Pi_{i=1}^s Sp_{2r_i}(F),$$

where  $r_i$  is the number  $H_{k_i}$ 's in the decomposition of  $M_{l_i}$ . Since  $M$  has decomposition (3.4),  $M_-$  has decomposition

$$M_- = F[t]/(t^{k_1}) \oplus \cdots \oplus F[t]/(t^{k_n}). \quad (7.2)$$

Let  $T$  be a maximal torus of  $H$ , we may take  $T$  such that  $T$  preserves  $M_{\pm}$  and preserves each component in (7.2). Then  $T = G_{\mathbf{m}}^n$ , where the  $i$ -th  $G_m$

acts on the  $i$ -th component in (7.2). We have  $T_{\mathbf{A}} = I_F^n$ , where  $I_F$  denotes the idele group of  $F$ . Let  $T'_{\mathbf{A}}$  be the subset of  $T_{\mathbf{A}}$  formed by

$$T'_{\mathbf{A}} = \{(t_1, \dots, t_n) \mid |t_1|_{\mathbf{A}} \geq \dots \geq |t_{r_1}|_{\mathbf{A}} \geq 1, \dots, |t_{n-r_s+1}|_{\mathbf{A}} \geq \dots \geq |t_n|_{\mathbf{A}} \geq 1\}.$$

Since  $N$  is unipotent, the space  $N(F) \backslash N(\mathbf{A})$  is compact. By the reduction theory for the semi-simple group  $H$ , there exists a compact  $C \subset Sp(M, t)_{\mathbf{A}}$  such that

$$Sp(M, t)_{\mathbf{A}} = Sp(M, t)_F T'_{\mathbf{A}} C. \quad (7.3)$$

Since  $T$  preserves  $M_+$ , we may regard  $T_{\mathbf{A}}$  as a subgroup of  $\widehat{Sp}(M, t)_{\mathbf{A}}$ , the decomposition (7.3) implies that

$$\widehat{Sp}(M, t)_{\mathbf{A}} = Sp(M, t)_F T'_{\mathbf{A}} C. \quad (7.4)$$

for some compact subset  $C \subset \widehat{Sp}(M, t)_{\mathbf{A}}$ .

As in [7], for each  $\tau \in \mathbb{R}_{>0}$ , we let  $a_{\tau} \in I_F$  denote the idele such that  $(a_{\tau})_v = \tau$  for each infinite place  $v$  and  $(a_{\tau})_v = 1$  for each finite place. We let  $\Theta(T)$  denote the set of all  $(a_{\tau_1}, \dots, a_{\tau_n})$ , and set  $\Theta(T)' = \Theta(T) \cap T'_{\mathbf{A}}$ .

**Lemma 7.1** *If  $\hat{E}$  is a positive tempered measure on  $X_{\mathbf{A}} = (M_- \otimes V)_{\mathbf{A}}$ , and is a sum of positive measures  $\hat{\mu}_i$  supported on  $U(i)_{\mathbf{A}}$  ( $i \in S_t^2(M_-)$ ), and is  $T_F$ -invariant, and there is a place  $v$  of  $F$  and a subgroup  $G'_v$  of  $G^q(F_v[[t]])$  that acts transitively on  $U(i)_v$  such that  $\hat{E}$  is invariant under  $G'_v$ . Then the function  $S \mapsto \hat{E}(S\phi)$  is bounded on  $T'_{\mathbf{A}}$ , uniformly for  $\phi$  in a compact subset in  $\mathcal{S}(X_{\mathbf{A}})$ .*

This lemma is a generalization of Lemma 23 in [7]. Our proof below closely follows that of [7].

*Proof.* Let  $e_1, \dots, e_n$  be a quasi-basis of  $M_-$ :

$$M_- = F[t]/(t^{k_1})e_1 \oplus \dots \oplus F[t]/(t^{k_n})e_n.$$

Then  $e_i \otimes e_j + e_j \otimes e_i$  is a quasi-basis of  $\text{St}^2(M_-)$ . For each  $\alpha \in \{0, 1, \dots, n\}$ , let  $S_t^2(M_-)^{(\alpha)}$  be the set that consists of elements

$$\sum_{i,j > \alpha} k_{ij}(e_i \otimes e_j + e_j \otimes e_i)$$

such that

$$k_{\alpha+1,j}(e_{\alpha+1} \otimes e_j + e_j \otimes e_{\alpha+1}) \neq 0$$

for at least one  $j \geq \alpha + 1$ . We set

$$S_t^2(M_-)^{(n)} = \{0\}$$

by convention. It is clear that  $S_t^2(M_-)$  is a disjoint union of  $S_t^2(M_-)^{(\alpha)}$ . Let  $\hat{E}_\alpha$  be the sum of  $\hat{\mu}_i$  for  $i \in S_t^2(M_-)^{(\alpha)}$ . It is clear that

$$\hat{E} = \hat{E}_0 + \hat{E}_1 + \cdots + \hat{E}_n$$

and  $\hat{E}_\alpha$  satisfies all the conditions in the lemma. it is enough to prove the result for each  $\hat{E}_\alpha$ .

Now we fix  $0 \leq \alpha \leq n$ . Let  $q$  be the constant which is 1 if  $v$  is infinite and is equal to the cardinality of the residue field if  $v$  is a finite place. As in [7], there is a compact subset  $C \subset I_F$  (where  $I_F$  is the idele group of  $F$ ), such that every  $t \in I_F$  with  $1 \leq |t| \leq S$  (where  $S$  is a fixed constant) can be written as  $rc$  with  $r \in F$ ,  $c \in C$ . We denote  $C^n$  the subset of  $T_{\mathbf{A}}$  formed by elements  $(c_1, \dots, c_n)$  with all  $c_i \in C$ , and let

$$\Theta'_\alpha = \{(a_{\tau_1}, \dots, a_{\tau_n}) \mid \tau_1 = \cdots = \tau_{\alpha+1} \geq \cdots \geq \tau_n \geq 1\}.$$

We will apply Lemma 6 of [7] to the space  $X_F$ . We consider  $X_F$  as a product space  $\prod_{i=1}^{n-\alpha} X_F^{(i)}$ , where  $X_F^{(1)}$  is the  $F[[t]]$ -submodule generated by  $e_1 \otimes V, \dots, e_{\alpha+1} \otimes V$ , and for  $n - \alpha \geq i \geq 2$ ,  $X_F^{(i)}$  is  $F[[t]]e_{i+\alpha}$ ,  $Y_F \subset M_- \otimes_{F[[t]]} M_-$  is the submodule over  $F[[t]]$  spanned by  $e_i \otimes e_j$  with  $j \leq \alpha + 1$ . And  $p : X_F \rightarrow Y_F$  is given by

$$p\left(\sum_{i=1}^{j=1} e_i \otimes v_i\right) = \sum_{i=1}^n \sum_{j=1}^{\alpha+1} (v_i, v_j) e_i \otimes e_j.$$

Apply Lemma 6 of [7], there is  $\phi_0 \in S(X_{\mathbf{A}})$  such that

$$|((\theta c) \cdot \phi)(x)| \leq \phi_0(x) \tag{7.5}$$

for all  $x \in X_{\mathbf{A}}$  with  $T(x) \in S_t(M_-)^{(\alpha)}$ , all  $\theta \in \Theta'_\alpha$ ,  $c \in C^n$ , and  $\phi \in C_0$ .

It can be proved ([7] page 71) that each element  $t = (t_1, \dots, t_n) \in T'_{\mathbf{A}}$  can be written as

$$t = ry\theta c$$

where  $r \in T_F$ ,  $y = (y_1, \dots, y_n) \in T_v$  with all  $|y_i|_v \geq 1$  and  $y_{\alpha+1} = \cdots = y_n = 1$ ,  $\theta \in \Theta'_\alpha$  and  $c \in C^n$ . Since  $\hat{E}_\alpha$  is invariant under  $T_F$ , we have

$$|\hat{E}_\alpha(t \cdot \phi)| \leq \hat{E}_\alpha(y \cdot \phi_0).$$

We may assume that  $\phi_0 = \phi_v \phi'$  where  $\phi_v \in \mathcal{S}(X_v)$ ,  $\phi' \in \mathcal{S}(\Pi'_{w \neq v} X_w)$ . By Lemma 22 [7], we have

$$\hat{E}_\alpha(y \cdot \phi_0) = \sum_{i \in St^2(M_-)^{(\alpha)}} c_i(\phi') \int_{U(i)_v} y \cdot \phi_v |\theta_i|_v.$$

From this, we obtain that

$$\hat{E}_\alpha(y \cdot \phi_0) = \Pi_{i=1}^n |y_i|_v^{k_1 + \dots + k_{i-1} + (n-m/2-i+2)k_i} \hat{E}_\alpha(\phi_0)$$

since  $k_i \leq k_j$  for  $i \leq j$ , and  $m > 6n + 2$ , we see that the exponent of  $|y_i|_v$  is  $\leq 0$ , and since  $|y_i|_v \geq 1$ , so have

$$\hat{E}_\alpha(y \cdot \phi_0) \leq \hat{E}_\alpha(\phi_0).$$

This proves the lemma.  $\square$

The following theorem is a generalization of Theorem 4 in [7].

**Theorem 7.2** *Suppose  $\dim V > 6n + 2$ . If there is a place  $v$  of  $F$  such that  $U(0)_v$  is not empty, and a subgroup  $G'_v$  of  $G^q(F_v[[t]])$  acts transitively on  $U(i)_v$  for every  $i \in S_t^2(M_-)$ . And if  $E'$  is a positive tempered measure on  $X_{\mathbf{A}}$  invariant under  $Sp(M, t)$  and  $G'_v$  and  $E' - \text{Et}$  is supported on the union of  $U(i)_{\mathbf{A}}$  for  $i \in S_t^2(M_-)$ . Then  $E' = \text{Et}$ .*

*Sketch of Proof.* Using Lemma 7.1 and 7.4, it is easy to see that the function  $\widehat{Sp}(M, t)_{\mathbf{A}} \rightarrow \mathbb{C}$  given by  $S \mapsto (E' - \text{Et})(S\phi)$  is bounded on  $\widehat{Sp}(M, t)_{\mathbf{A}}$  uniformly for  $\phi$  in every compact subset of  $\mathcal{S}(X_{\mathbf{A}})$ . The remainder of the proof is similar to that of Theorem 4 in [7].  $\square$

Now we can prove the main theorem of this work:

**Theorem 7.3** *Suppose  $\dim V > 6n + 2$  and  $V$  is anisotropic or  $\dim V - r > \frac{1}{2}\dim M + 1$ , where  $r$  is the dimension of a maximal isotropic subspace of  $V$ , then*

$$\text{Et} = \text{It}.$$

The condition on  $V$  in the Theorem is for the convergence of  $\text{It}$  (see 6.1). The proof uses the induction on  $\dim M$ . The induction assumption implies that  $E' = \text{It}$  satisfies the conditions of Theorem 7.2, therefore  $\text{Et} = \text{It}$ .

## 8 Corollaries of Siegel-Weil formula

In this section we prove a slightly more general form of the Siegel-Weil formula (Theorem 8.1) for snt-modules that will be used in part II. Let  $M, V$  be as in Section 3. Let

$$M = M_- \oplus M_+ \quad (8.1)$$

be a direct sum such that  $M_+ \in Gr(M, t)$ ,  $M_- \in Gr(M)$  but not necessarily in  $Gr(M, t)$ . Let  $X = M_- \otimes V$ . The space  $L^2(X_{\mathbf{A}})$  is a representation of the metaplectic group  $\widehat{Sp}_{2N}(\mathbf{A})$  ( $2N = \dim M \dim V$ ) with the usual theta functional

$$\theta : \mathcal{S}(X_{\mathbf{A}}) \rightarrow \mathbb{C}, \quad \phi \mapsto \theta(\phi) = \sum_{r \in X} \phi(r).$$

Recall the Eisenstein series (1.3), (1.7) for  $\phi \in \mathcal{S}(X_{\mathbf{A}})$  is given by

$$E(\phi) = \sum_{U \in Gr(M)} E(\phi, U) = \sum_{U \in Gr(M)} \int_{(\pi_-(U) \otimes V)_{\mathbf{A}}} \psi\left(\frac{1}{2}\langle x, \rho x \rangle\right) \phi(x) dx.$$

The  $t$ -Eisenstein series defined in (1.11) is a subseries given by

$$Et(\phi) = \sum_{U \in Gr(X, t)} E(\phi, U). \quad (8.2)$$

Though  $M_-$  is not an  $F[[t]]$ -submodule of  $M$ , it has  $F[[t]]$ -module structure via the isomorphism  $M_- = M/M_+$ . In the case that  $M_-$  is an  $F[[t]]$ -submodule, the two  $F[[t]]$ -module structures on  $U_-$  clearly coincide. The projection map

$$\pi_- : M \rightarrow M/M_+ = M_-$$

is an  $F[[t]]$ -module homomorphism. For each  $U \in Gr(M, t)$ ,  $\pi_-(U)$  is an  $F[[t]]$ -submodule of  $M_-$ . Denote  $Gr(M_-, t)$  the set of  $F[[t]]$ -submodules of  $M_-$ , so we have a map

$$P : Gr(M, t) \rightarrow Gr(M_-, t) : U \mapsto \pi_-(U). \quad (8.3)$$

For  $W \in Gr(M_-, t)$ , we set

$$Et_W(\phi) = \sum_{U \in Gr(M, t) : \pi_-(U) = W} E(\phi, U).$$



We recall the action formula of  $G^q(\mathbf{A}[[t]])$  on  $\mathcal{S}((M_- \otimes V)_{\mathbf{A}})$ . An element  $g \in G^q(\mathbf{A}[[t]])$  has the block decomposition

$$\begin{bmatrix} \alpha_g & \beta_g \\ \gamma_g & \delta_g \end{bmatrix}$$

with respect the decomposition

$$X_{\mathbf{A}} = (M_- \otimes V)_{\mathbf{A}} \oplus (M_+ \otimes V)_{\mathbf{A}}.$$

Since  $G^q(\mathbf{A}[[t]])$  preserves  $(M_+ \otimes V)_{\mathbf{A}}$ , so we have  $\gamma_g = 0$ . Since  $M_-$  is not an  $F[[t]]$ -submodule in general,  $\beta_g$  may not be 0. Then the action of  $g$  on  $\mathcal{S}((X \otimes V)_{\mathbf{A}})$  is given by

$$(g \cdot \phi)(x) = \psi\left(\frac{1}{2}\langle x\alpha_g, x\beta_g \rangle\right)\phi(x\alpha_g) \quad (8.4)$$

If  $g \in G^q(F[[t]])$  and  $\xi \in V$ , then

$$\frac{1}{2}\langle \xi\alpha_g, \xi\beta_g \rangle \in F,$$

and so  $\psi\left(\frac{1}{2}\langle \xi\alpha_g, \xi\beta_g \rangle\right) = 1$  and

$$(g \cdot \phi)(\xi) = \phi(\xi\alpha_g).$$

Therefore, we have

$$\theta(g \cdot \phi) = \theta(\phi), \quad \text{for } g \in G^q(F[[t]]). \quad (8.5)$$

The function  $\theta(g \cdot \phi)$  (as a function on  $G^q(\mathbf{A}[[t]])$ ) is actually a function on  $G^q(F[[t]]) \backslash G^q(\mathbf{A}[[t]])$ . Assume  $(V, (\cdot, \cdot))$  satisfies the conditions in Lemma 6.1, then we can form the convergent integral

$$\text{It}(\phi) \stackrel{\text{def}}{=} \int_{G^q(F[[t]]) \backslash G^q(\mathbf{A}[[t]])} \theta(g \cdot \phi) dg, \quad (8.6)$$

the convergence can be proved in the same way as Lemma 6.1.

We wish to write the integral (8.6) as a sum of orbital integrals. We introduce the set

$$\Omega = S^1 \times (M_- \otimes V)_{\mathbf{A}},$$

on which  $G^q(\mathbf{A}[[t]])$  acts as

$$(s, x) \cdot g = (s\psi\left(\frac{1}{2}\langle x\alpha_g, x\beta_g \rangle\right), x\alpha_g). \quad (8.7)$$

One can then check directly that (8.7) does define a group action; i.e., that

$$((s, x)g_1)g_2 = (s, x)g_1g_2, \quad g_1, g_2 \in G^q(\mathbf{A}[[t]]).$$

On the other hand, if

$$\phi \in \mathcal{S}((U_- \otimes V)_{\mathbf{A}}),$$

we can extend  $\phi$  to  $\Omega$  by

$$\varphi(s, x) = s\varphi(x), \quad x \in (U_- \otimes V)_{\mathbf{A}}, \quad s \in S^1.$$

And note that we can rewrite the integral (8.6) as

$$\int_{G^q(F[[t]]) \backslash G^q(\mathbf{A}[[t]])} \left( \sum_{\xi \in U_- \otimes V} \phi((1, \xi) \cdot g) \right) dg. \quad (8.8)$$

We note that the subset

$$(1, U_- \otimes V) \subset \Omega$$

is invariant under  $G^q(F[[t]])$ . We let  $\mathcal{O} \subset (1, U_- \otimes V)$  be a family of orbit representatives for the action of  $G^q(F[[t]])$ . Of course  $\mathcal{O}$  can be identified with a subset of  $U_- \otimes_F V$  ( $(1, \xi) \mapsto \xi$ ,  $\xi \in U_- \otimes V$ ), and as such, it is a family of orbit representatives for the action

$$\xi \mapsto \xi\alpha_g, \quad g \in G^q(F[[t]])$$

of  $G^q(F[[t]])$  on  $U_- \otimes V$ . We may rewrite (8.8) as

$$\int_{G^q(F[[t]]) \backslash G^q(\mathbf{A}[[t]])} \sum_{\xi \in \mathcal{O}} \sum_{\tau \in G_{\xi} \backslash G^q(F[[t]])} \varphi((1, \xi)\tau g) dg. \quad (8.9)$$

Using the fact that  $G_{\xi, \mathbf{A}}$  is unimodular, we can prove (8.9) is equal to

$$\text{It}(\phi) = \sum_{\xi \in \mathcal{O}} \text{vol}(G_{\xi} \backslash G_{\xi, \mathbf{A}}) \int_{G_{\xi, \mathbf{A}} \backslash G^q(\mathbf{A}[[t]])} \phi((1, \xi)g_2) dg_2, \quad (8.10)$$

by a similar argument as in [7] (page 16). We have thereby expressed the  $\text{It}(\phi)$  as a sum of orbital integrals. For each orbit  $\mathcal{O}$ , we have a corresponding  $W \in \text{Gr}(M_-, t)$  (see Theorem 5.8). Let  $\text{It}_W(\phi)$  be the subseries of (8.10) that is over all  $\xi$  corresponding to  $W$ . Then we have

$$\text{It}(\phi) = \sum_{W \in \text{Gr}(M_-, t)} \text{It}_W(\phi).$$

**Theorem 8.1** *Suppose  $\dim V > 6n + 2$  and  $(V, (\cdot, \cdot))$  satisfies the condition in Lemma 6.1. Let  $M = M_- \oplus M_+$  be a decomposition such that  $M_- \in \text{Gr}(M)$  and  $M_+ \in \text{Gr}(M, t)$ . Put  $X = M_- \otimes V$ . We can define tempered distributions  $\text{Et}$  and  $\text{It}$  on  $X_{\mathbf{A}}$  as in (8.2) and (8.6). Then*

$$\text{Et} = \text{It}. \quad (8.11)$$

And for each  $W \in \text{Gr}(M_-, t)$ , we have

$$\text{Et}_W = \text{It}_W. \quad (8.12)$$

Notice that we didn't assume  $M_- \in \text{Gr}(M, t)$ , but this theorem can be reduced to Theorem 7.3. Let

$$M = \bar{M}_- \oplus \bar{M}_+$$

be a decomposition of  $t$ -Lagrangian subspaces. Both  $L^2((M_- \otimes V)_{\mathbf{A}})$  and  $L^2((\bar{M}_- \otimes V)_{\mathbf{A}})$  are models of the Weil representation of  $\widehat{Sp}_{2N}(\mathbf{A})$ . We shall define an intertwining operator

$$T : L^2((M_- \otimes V)_{\mathbf{A}}) \rightarrow L^2((\bar{M}_- \otimes V)_{\mathbf{A}})$$

such that  $T$  sends  $\mathcal{S}((M_- \otimes V)_{\mathbf{A}})$  to  $\mathcal{S}((\bar{M}_- \otimes V)_{\mathbf{A}})$ , and

$$\text{Et}(\phi) = \text{Et}(T\phi), \quad \text{It}(\phi) = \text{It}(T\phi).$$

By Theorem 7.3,  $\text{Et}(T\phi) = \text{It}(T\phi)$ , so we have  $\text{Et}(\phi) = \text{It}(\phi)$ . The more detailed proof follows.

*Proof.* Recall for the symplectic space  $M \otimes V$ , we have the associated Heisenberg group

$$H = (M \otimes V)_{\mathbf{A}} \times S^1$$

with the product given by

$$(a_1, s_1)(a_2, s_2) = (a_1 + a_2, s_1 s_2 \psi(\frac{1}{2} \langle a_1, a_2 \rangle)).$$

Since  $M_+$  is a Lagrangian subspace,  $A \stackrel{\text{def}}{=} (M_+ \otimes V)_{\mathbf{A}} \times S^1$  is a maximal abelian subgroup of  $H$ . And

$$\chi : (M_+ \otimes V)_{\mathbf{A}} \times S^1 \rightarrow S^1, \quad (a, s) \mapsto s$$

is a 1-dimensional representation. The induced representation

$$\text{Ind}_A^H \chi$$

consists of functions  $f$  on  $H$  such that  $f(ax) = \chi(a)f(x)$  for all  $a \in A$  and the restriction  $f$  on  $(M_- \otimes V)_{\mathbf{A}}$  is in  $L^2((M_- \otimes V)_{\mathbf{A}})$ . Let  $M = \bar{M}_- \oplus \bar{M}_+$  be a direct sum of  $t$ -Lagrangian subspaces. Similarly, for the maximal abelian subgroup  $(\bar{M}_- \otimes V)_{\mathbf{A}} \times S^1$ , and the character

$$\bar{\chi} : (\bar{M}_- \otimes V)_{\mathbf{A}} \times S^1 \rightarrow S^1, \quad (\bar{a}, s) \mapsto s$$

we have the induced representation

$$Ind_{\bar{A}}^H \bar{\chi}.$$

The representation space  $Ind_A^H \chi$  ( $Ind_{\bar{A}}^H \bar{\chi}$ , resp.) can be identified with  $L^2((M_- \otimes V)_{\mathbf{A}})$  ( $L^2((\bar{M}_- \otimes V)_{\mathbf{A}})$ , resp.) by the restricting a function on  $H$  to  $(\bar{M}_- \otimes V)_{\mathbf{A}}$  ( $(\bar{M}_- \otimes V)_{\mathbf{A}}$ , reps.). The space of smooth vectors are  $\mathcal{S}(M_- \otimes V)_{\mathbf{A}}$  and  $\mathcal{S}((\bar{M}_- \otimes V)_{\mathbf{A}})$ , respectively. Let  $h \in Sp(M)$  be an element such that

$$M_- h = \bar{M}_-, \quad M_+ h = \bar{M}_+.$$

We define

$$T : \mathcal{S}(M_- \otimes V)_{\mathbf{A}} \rightarrow \mathcal{S}((\bar{M}_- \otimes V)_{\mathbf{A}})$$

by

$$(Tf)(x) = (h \cdot f)(xh^{-1}).$$

It is easy to check that  $T$  is an isomorphism of  $H$ -representations. There are two actions of  $\widehat{Sp}_{2N}(\mathbf{A})$  on  $\mathcal{S}((\bar{M}_- \otimes V)_{\mathbf{A}})$  (or on  $L^2(\bar{M}_- \otimes V)_{\mathbf{A}}$ ): the Weil representation action which we denote by  $\pi(g)$ , and  $\pi'(g) = T\pi(g)T^{-1}$  (where  $\pi(g)$  denote the Weil representation action on  $\mathcal{S}(M_- \otimes V)_{\mathbf{A}}$ ). They both satisfy that, for every  $\alpha \in H$  :

$$\pi(g^{-1})\pi(\alpha)\pi(g) = \pi(\alpha \cdot g)$$

$$\pi'(g^{-1})\pi(\alpha)\pi'(g) = \pi(\alpha \cdot g)$$

So  $\pi'(g) = c_g \pi(g)$  for some scalar  $c_g$ . It is clear that  $c_{g_1 g_2} = c_{g_1} c_{g_2}$ . Since  $Sp_{2N}(F)$  is a perfect group, we have  $c_g = 1$  for all  $g \in Sp_{2N}(F)$ . This implies that  $c_g = 1$  for all  $g \in \widehat{Sp}_{2N}(\mathbf{A})$ . It then follows that  $T$  is an isomorphism of the Weil representations. We define as usual the theta function functionals

$$\theta : \mathcal{S}((M_- \otimes V)_{\mathbf{A}}) \rightarrow \mathbb{C}, \quad \bar{\theta} : \mathcal{S}((\bar{M}_- \otimes V)_{\mathbf{A}}) \rightarrow \mathbb{C}.$$

We have

$$\bar{\theta}(Tf) = \sum_{r \in \bar{M}_-} (Tf)(r) = \sum_{r \in \bar{M}_-} (h \cdot f)(rh^{-1}) = \sum_{r \in M_-} (h \cdot f)(r) = \theta(h \cdot f) = \theta(f).$$

And

$$\begin{aligned}
\text{It}(Tf) &= \int_{G^q(F[[t]] \backslash G^q(\mathbf{A}[[t]])} \theta(g \cdot Tf) dg \\
&= \int_{G^q(F[[t]] \backslash G^q(\mathbf{A}[[t]])} \bar{\theta}(T(g \cdot f)) dg \\
&= \int_{G^q(F[[t]] \backslash G^q(\mathbf{A}[[t]])} \theta((g \cdot f)) dg \\
&= \text{It}(f).
\end{aligned}$$

Let  $P \subset Sp(M)$  be the parabolic subgroup that consists of all  $g \in Sp(M)$  such that  $M_+g = M_+$ , similarly, let  $\bar{P} \subset Sp(M)$  be the parabolic subgroup that consists of all  $g \in Sp(M)$  such that  $\bar{M}_+g = \bar{M}_+$ . And let  $S$  denote the set of all  $g \in P \backslash Sp(M)$  such that  $M_+g$  is a  $t$ -Lagrangian subspace. Similarly let  $\bar{S}$  denote the set of all  $g \in \bar{P} \backslash Sp(M)$  such that  $\bar{M}_+g$  is a  $t$ -Lagrangian subspace. Since  $M_+h = \bar{M}_+$ , the map  $L_h : \bar{S} \rightarrow S, g \mapsto hg$  is a bijection.

$$\begin{aligned}
\text{Et}(Tf) &= \sum_{g \in \bar{S}} (g \cdot Tf)(0) \\
&= \sum_{g \in \bar{S}} (Tgf)(0) \\
&= \sum_{g \in \bar{S}} (hgf)(0) \\
&= \sum_{g \in S} (gf)(0) \\
&= \text{Et}(f).
\end{aligned}$$

This proves  $\text{Et}(\phi) = \text{It}(\phi)$ . To prove  $\text{Et}_W(\phi) = \text{It}_W(\phi)$  for every  $W \in Gr(M_-, t)$ , we first note for every  $W \in Gr(M_-, t)$ ,  $W + M_+$  is an  $F[[t]]$ -submodule of  $M$ . Let

$$W^\perp = \{v \in M_+ \mid \langle v, W \rangle = 0\}.$$

It is easy to see that  $W^\perp$  is the radical of the restriction of  $\langle, \rangle$  on  $W + M_+$ . Therefore

$$M_W \stackrel{\text{def}}{=} (W + M_+)/W^\perp$$

is an snt-module. And

$$M_W = W \oplus M_+/W^\perp$$

is a decomposition into Lagrangian subspaces, and  $M_+/W^\perp \in Gr(M_W, t)$ . We consider  $\mathcal{S}((W \otimes V)_{\mathbf{A}})$ , and apply (8.11) to the pair  $(Sp(M_W, t), G)$ , we have

$$\sum_{W' \in Gr(M_-, t): W' \subset W} \text{Et}_{W'}(\phi) = \sum_{W' \in Gr(M_-, t): W' \subset W} \text{It}_{W'}(\phi).$$

The above equality holds for all  $W \in Gr(M_-, t)$ , which implies  $\text{Et}_W(\phi) = \text{It}_W(\phi)$ .  $\square$

## References

- [1] H. Garland, Certain Eisenstein series on loop groups: convergence and the constant term, *Algebraic Groups and Arithmetic* Tata Inst. Fund. Res. Mumbai (2004), 275-319.
- [2] H. Garland, Absolute convergence of Eisenstein series on loop groups, *Duke Math J.* Vol. 135, No. 2 (2006), 203-260.
- [3] H. Garland, Y.Zhu, On the Siegel-Weil theorem for loop groups (II), preprint, 2008.
- [4] D.G. James, On Witt's theorem for unimodular, quadratic forms II, *Pacific Jour. of Math.* 33, 1970.
- [5] S.S. Kudla, S. Rallis, On the Weil-Siegel formula, *J. reine angew. Math.* 387 (1988), 1-68.
- [6] A. Weil, Sur certaines groupes d'opérateurs unitaires, *Acta Math.* 111 (1964), 143-211.
- [7] A. Weil, Sur la formule de Siegel dans la theorie des groupes calssiques, *Acta Math.* 113 (1965), 1-88.
- [8] Y. Zhu, Theta functions ans Weil representations of loop symplectic groups, *Duke Math. J.*, vol 143 (2208), no. 1, 17-39.

Dept of Math, Yale University, New Haven, CT 06520-8283. hgarland@math.yale.edu

Dept of Math, Hong Kong University of Science and Tecgnology. mazhu@ust.hk